



<http://ir4.rutgers.edu/Ornamental/ornamentalSummaryReports.cfm>

**IR-4 Ornamental Horticulture Program  
Fusarium Efficacy: A Literature Review**

*Fusarium avenaceum*  
*Fusarium commune*  
*Fusarium oxysporum*  
*Fusarium solani*

**Authors: Ely Vea and Cristi Palmer  
Date: June 11, 2013**

## Table of Contents

Table of Contents .....	2
Table of Tables .....	3
Abstract .....	5
Introduction.....	5
Materials and Methods.....	5
Results.....	9
Comparative Efficacy on <i>Fusarium avenaceum</i> .....	9
Comparative Efficacy on <i>Fusarium commune</i> .....	9
Comparative Efficacy on <i>Fusarium oxysporum</i> .....	15
Comparative Efficacy on <i>Fusarium solani</i> .....	28
Efficacy Summary by Product/Active Ingredient.....	32
Acibenzolar.....	32
Azoxystrobin.....	32
Boscalid + Pyraclostrobin.....	32
Caprylic Acid.....	32
Cyprodinil + Fludioxonil.....	32
Fludioxonil.....	32
Fludioxonil + Mefenoxam.....	33
Fluoxastrobin.....	33
Metconazole.....	33
Prothioconazole.....	33
Pyraclostrobin.....	33
SP2169.....	33
Thiamethoxam.....	33
Tebuconazole.....	33
Thiophanate methyl.....	34
Trichoderma harzianum & T. virens.....	34
Trifloxystrobin.....	34
Triflumizole.....	34
Triticonazole.....	34
Phytotoxicity.....	34
Appendix 1: Contributing Researchers .....	44
Appendix 2: Submitted Data Reports .....	45

## Table of Tables

Table 1.	List of Products and Rates Tested on Ornamentals from 2001 to 2012. ....	6
Table 2.	List of Products and Rates Tested on Vegetables from 2002 to 2011. ....	8
Table 3.	* Efficacy on Fusarium Crown and Stem Rot ( <i>Fusarium avenaceum</i> ) on Lisianthus ( <i>Eustoma grandiflorum</i> ) ‘Maurine Blue’, McGovern, FL, 2001. ....	9
Table 4.	Efficacy on Damping-off and Root Rot Caused by <i>Fusarium commune</i> on Douglas Fir ( <i>Pseudotsuga menziesii</i> ), Chastagner WA, 2010. ....	10
Table 5.	Efficacy on Damping-off and Root Rot Caused by <i>Fusarium commune</i> on Douglas Fir ( <i>Pseudotsuga menziesii</i> ), Chastagner WA, 2011, Trial 1. ....	11
Table 6.	Efficacy on Damping-off and Root Rot Caused by <i>Fusarium commune</i> on Douglas Fir ( <i>Pseudotsuga menziesii</i> ), Chastagner WA, 2011, Trial 2. ....	11
Table 7.	Efficacy on Damping-off and Root Rot Caused by <i>Fusarium commune</i> on Douglas Fir ( <i>Pseudotsuga menziesii</i> ), Chastagner WA, 2012. ....	13
Table 8.	Efficacy on Damping-off and Root Rot Caused by <i>Fusarium commune</i> on Bulbous Iris ( <i>Iris x hollandica</i> ) ‘Blue Diamond’, Chastagner WA, 2012. ....	14
Table 9.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> ) on Lisianthus ( <i>Eustoma grandiflorum</i> ) ‘Maurine Blue’, McGovern, FL, 2001. ....	15
Table 10.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> ) on Lisianthus ( <i>Eustoma grandiflorum</i> ) ‘Maurine Blue’, McGovern, FL, 2001. ....	15
Table 11.	Efficacy on Fusarium Root Rot ( <i>Fusarium oxysporum</i> ) on Larkspur ( <i>Delphinium sp.</i> ), Kirk, MI, 2010. ....	17
Table 12.	Efficacy on Fusarium wilt ( <i>F. oxysporum</i> ) on Lisianthus ( <i>Eustoma grandiflora</i> ), Chase, CA, 2011. ....	18
Table 13.	Efficacy on Stem Based Necrosis Caused By <i>Fusarium oxysporum</i> on Corn Flag ( <i>Gladiolus sp.</i> ) ‘Sun-Kissed’, Kirk, MI, 2012, Trial 1. ....	20
Table 14.	Efficacy on Stem Based Necrosis Caused By <i>Fusarium oxysporum</i> on Corn Flag ( <i>Gladiolus sp.</i> ) ‘Sun-Kissed’, Kirk, MI, 2012, Trial 2. ....	21
Table 15.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Egel, IN, 2007. ....	22
Table 16.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Egel, IN, 2008. ....	22
Table 17.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Langston, GA, 2008. ....	23
Table 18.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Sugar Baby’, Everts, MD, 2008. ....	23
Table 19.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Sugar Baby’, Everts, DE, 2008. ....	24
Table 20.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Sugar Baby’, Everts, MD, 2008. ....	25
Table 21.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Sugar Baby’, Everts, MD, 2009. ....	26
Table 22.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Imagination’, Egel, IN, 2009. ....	26
Table 23.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Langston, GA, 2010. ....	27

Table 24.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Langston, GA, 2011, Trial 1. ....	27
Table 25.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Langston, GA, 2011, Trial 2. ....	28
Table 26.	* Efficacy on Fusarium Wilt ( <i>Fusarium oxysporum</i> f. sp. <i>niveum</i> ) on Watermelon ( <i>Citrullus lanatus</i> ) ‘Black Diamond’, Langston, GA, 2011, Trial 3. ....	28
Table 27.	* Efficacy on Fusarium Tuber Rot ( <i>Fusarium solani</i> ) on Caladium ( <i>Caladium x hortulanum</i> ) ‘Florida Cardinal’, McGovern, FL, 2001.....	29
Table 28.	Efficacy on Stem Root Rot Caused by <i>Fusarium solani</i> on <i>Dracaena deremensis</i> ‘Janet Craig’, Palmateer FL, 2011. ....	30
Table 29.	* Efficacy on Fusarium Root Rot ( <i>Fusarium solani</i> f. sp. <i>phaseoli</i> ) on Snap Bean ( <i>Phaseolus vulgaris</i> ) ‘Hi-Style’, Kirk, MI, 2002.....	31
Table 30.	Summary of product efficacy by pathogen and crop. ....	35

## Abstract

From 2001 to 2012, numerous products representing 27 active ingredients were evaluated in greenhouse and field trials as soil drench, soil incorporation, foliar, in-furrow, drip irrigation or tuber soak applications against several *Fusarium* species causing rots (crown, stem and tuber rots) and wilt on ornamentals, and wilt and root rot on vegetables (Table 1, Table 2). *Fusarium* species tested included: *F. avenaceum*, *F. commune*, *F. oxysporum* and *F. solani*. Most trials were conducted on *F. oxysporum* on larkspur, lisianthus and watermelon. Although there were insufficient data for definitive conclusions, several relatively new products showed promising, though inconsistent, efficacy comparable to the standards. These include acibenzolar, Heritage (azoxystrobin), Compass (trifloxystrobin), Hurricane (fludioxonil + mefenoxam), Insignia (pyraclostrobin), SP2169, Tourney (metconazole) and Trinity (triticonazole). BW240/RootShield Plus (*Trichoderma harzianum* & *T. virens*), CG100 (caprylic acid), Pageant (boscalid + pyraclostrobin) and Palladium (cyprodinil + fludioxonil) provided no to mediocre efficacy. Proline (prothioconazole) provided consistently good control of *F. oxysporum* in watermelon trials. The established standards 3336 and Medallion generally provided inconsistent efficacy while Terraguard was effective in one trial.

## Introduction

In 2010, IR-4 initiated a high priority project to determine efficacy of several fungicides on *Fusarium* species and obtain data supporting current and future registrations on ornamentals. There are many different species of *Fusarium* causing ornamental diseases and an extensive project may be required to generate sufficient efficacy data. We reviewed available ornamental and vegetable trials published in Biological & Cultural Tests, Fungicide & Nematicide Tests and Plant Disease Management Reports to check efficacy of experimental and registered fungicides on *Fusarium* species. This report is a brief summary of available data from 14 ornamental and 13 vegetable trial reports. The source of report is included under each data table. Ten trials from the IR-4 project are included in this report. Additional data will be added when received from researchers.

## Materials and Methods

From 2001 to 2012, numerous products representing 27 active ingredients were evaluated in greenhouse and field trials as soil drench, soil incorporation, foliar, in-furrow, drip irrigation or tuber soak applications against several *Fusarium* species causing rots (crown, stem and tuber rots) and wilt on ornamentals, and wilt and root rot on vegetables (Table 1, Table 2). *Fusarium* species tested included: *F. avenaceum*, *F. communi*, *F. oxysporum* and *F. solani*. Most trials were conducted on *F. oxysporum* on larkspur, lisianthus and watermelon. In greenhouse studies, treatments were generally applied as soil drench either a few days before *Fusarium* inoculation or immediately after inoculation and reapplied biweekly. In field trials, treatments were generally applied as soil drench immediately after transplanting, through drip irrigation several times during the growing season, in-furrow at planting, tuber soak application or foliar spray for control of natural *Fusarium* infestations. Researchers used a minimum of four replications. Disease severity and incidence were recorded at various intervals after initial application. Phytotoxicity or lack of it was generally noted in the reports. Seven researchers were involved in the testing (Appendix 1).

Products were supplied by their respective manufacturers.

For IR-4 testing, the following protocols were used: 10-016, 11-010 and 12-016. Please visit <http://ir4.rutgers.edu/ornamental/OrnamentalDrafts.cfm> to view and download these protocols.

**Table 1. List of Products and Rates Tested on Ornamentals from 2001 to 2012.**

Active Ingredient(s)	Trade Name(s)		Manufacturer	Rate(s) Tested		# Trials
	Food Use	Orn.Hort./ Turf Use				
Acibenzolar	Actigard		Syngenta	Corm Dip	0.125 oz per 100 gal	1
					0.25 oz per 100 gal	1
					0.5 oz per 100 gal	1
				Drench	0.125 oz per 100 gal	5
					0.25 oz per 100 gal	
				Foliar	0.5 oz per 100 gal	1
0.75 oz per 100 gal	1					
Soil incorp.	0.08 g per 1000 ft <sup>2</sup>	2				
Azoxystrobin	Abound, Amistar, Quadris	Heritage	Syngenta	Corm Dip	0.45 oz per 100 gal	1
					0.9 oz per 100 gal	1
					1.8 oz per 100 gal	1
				Drench	0.9 oz per 100 gal	6
					1.8 oz per 100 gal	
					4 oz per 100 gal	
					8.7 oz per 100 gal	
				Foliar	16 oz per 100 gal	1
1.8 oz per 100 gal						
Sprench	2.7 oz per 100 gal	1				
	4 oz per 100 gal	1				
Soil incorp.	0.4 oz per 1000 ft <sup>2</sup>	2				
Boscalid + Pyraclostrobin	Pristine	Pageant	BASF	Drench	8 oz per 100 gal 12 oz per 100 gal	3
Caprylic acid	CG100	CG100	Summerdale	Corm Dip	38.4 fl oz per 100 gal	1
				Drench	9.6 fl oz per 100 gal	4
					0.6 pt per 100 gal 0.8 pt per 100 gal	
				Foliar	38.4 fl oz per 100 gal	1
				Sprench	0.6 pt per 100 gal	1
Soil incorp.	26 ml per 1000 ft <sup>2</sup>	2				
Copper sulfate pentahydrate		Phyton 27	Phyton	Drench	25 oz per 100 gal	1
Chlorothalonil + Thiophanate-methyl		Spectro	Cleary	Drench	12 oz per 100 gal 24 oz per 100 gal	2
Cyprodinil + fludioxonil	Switch	Palladium	Syngenta	Corm Dip	6 fl oz per 100 gal	1
				Drench	6 oz per 100 gal	3
					4 oz per 100 gal	
				Foliar	6 fl oz per 100 gal	1
				Sprench	4 oz per 100 gal	1
Soil incorp.	1 oz per 1000 ft <sup>2</sup>	2				
Fluazinam	Omega	Fluazinam	Syngenta	Drench	6.4 fl oz per 100 gal	1
Fludioxonil	Cannonball , Scholar	Medallion	Syngenta	Corm Dip	8 oz per 100 gal	1
				Drench	2 oz per 100 gal	6
				Foliar	4 oz per 100 gal	1
				Sprench	2 oz per 100 gal	1

Active Ingredient(s)	Trade Name(s)		Manufacturer	Rate(s) Tested		# Trials
	Food Use	Orn.Hort./ Turf Use				
Fludioxonil + Mefenoxam		Hurricane	Syngenta	Drench	0.75 oz per 100 gal 1.5 oz per 100 gal 12 oz per 100 gal	4
				Spray	1.5 oz per 100 gal	1
Fluoxastrobin	Evito	Disarm	Arysta, OHP	Corm Dip	4 fl oz per 100 gal	1
				Foliar	4 fl oz per 100 gal	1
Hydrogen peroxide, peroxyacetic & octanoic acids		X-3	Phyton	Drench	1:500 dilution	1
				Corm Dip	2 fl oz per 100 gal 4 fl oz per 100 gal	1
Metconazole	Caramba, Quash, V-10116	Tourney	BASF, Valent	Drench	1 oz per 100 gal 2 oz per 100 gal	4
				Foliar	1 fl oz per 100 gal 2 fl oz per 100 gal	1
				Sprenc	1 oz per 100 gal 2 oz per 100 gal	1
				Soil incorp.	2.8 g per 1000 ft <sup>2</sup> 5.6 g per 1000 ft <sup>2</sup>	2
Propiconazole	Orbit, Tilt	Banner Maxx	Syngenta	Drench	5 fl oz per 100 gal	1
Pyraclostrobin	Cabrio, Headline	Insignia	BASF	Drench	3.1 fl oz per 100 gal 6.1 fl oz per 100 gal	1
SP2169		SP2169	SePro	Corm Dip	12.3 fl oz per 100 gal	1
				Drench	12.3 fl oz per 100 gal	2
				Foliar	12.3 fl oz per 100 gal	1
				Soil incorp.	8.35 ml per 1000 ft <sup>2</sup>	2
Tebuconazole	Folicur	Torque	Bayer, Cleary	Drench	6 fl oz per 100 gal	1
				Soil incorp.	20.4 ml per 1000 ft <sup>2</sup>	2
Thiamethoxam	Actara, Platinum	Flagship 0.22 G	Syngenta	Soil incorp.	44 oz per 1000 ft <sup>2</sup>	2
		Flagship 25WG		Soil incorp.	0.55 oz per 1000 ft <sup>2</sup>	2
Triflumizole	Procure	Terraguard	Chemtura	Drench	6 oz per 100 gal	1
Thiophanate methyl	Topsin	3336	UPI, Cleary	Corm Dip	16 oz per 100 gal	1
				Drench	16 oz per 100 gal 24 oz per 100 gal	5
				Foliar	16 fl oz per 100 gal	1
				Sprenc	12 fl oz per 100 gal	1
<i>Trichoderma harzianum</i> & <i>T. virens</i>	RootShield Plus	RootShield Plus, BW240	BioWorks	Corm Dip	80 oz per 100 gal	1
				Drench	3 oz per 100 gal 6 oz per 100 gal 8 oz per 100 gal	4
				Foliar	8 oz per 100 gal	1
				Sprenc	6 oz per 100 gal	1
				Soil incorp.	10 oz per 1000 ft <sup>2</sup>	2
Trifloxystrobin	Flint, Gem	Compass	Bayer	Drench	0.5 oz per 100 gal 1 oz per 100 gal 4 oz per 100 gal	4
				Sprenc	2 oz per 100 gal	1
Triticonazole	Charter	Trinity	BASF	Drench	6 fl oz per 100 gal 8 fl oz per 100 gal	4
				Foliar	8 fl oz per 100 gal 12 fl oz per 100 gal	1

Active Ingredient(s)	Trade Name(s)		Manufacturer	Rate(s) Tested		# Trials
	Food Use	Orn.Hort./ Turf Use		Sprench	6 fl oz per 100 gal 8 fl oz per 100 gal	
					Soil incorp.	

**Table 2. List of Products and Rates Tested on Vegetables from 2002 to 2011.**

Active Ingredient(s)	Trade Name(s)		Manufacturer	Rate(s) Tested		# Trials
	Food Use	Orn.Hort./ Turf Use		Drench	0.33 oz per acre 0.25 oz per 100 gal 0.5 oz per 100 gal 0.75 oz per 100 gal	
					Drip	
Acibenzolar	Actigard		Syngenta	Foliar	0.75 oz per acre	1
				Drench	15.4 fl oz per 100 gal	9
				In-furrow	0.8 pt per 1000 ft	1
Azoxystrobin	Abound, Amistar, Quadris	Heritage	Syngenta	Foliar	15.4 fl oz per acre	1
				Drench	2 qt per 100 gal 4 qt per 100 gal	1
				In-furrow	1.5 pt per 1000 ft	1
<i>Bacillus subtilis</i>	Serenade	Rhapsody	AgraQuest	Drench	0.5 lb per acre 16 oz per 100 gal	5
Chlorothalonil	Bravo	Daconil	Syngenta	Drench	3 ml per sq m 6 ml per sq m	2
Fludioxonil	Cannonball , Scholar	Medallion	Syngenta	Drench	4 oz per 100 gal 8 oz per 100 gal	5
Hymexazol	Tachigaren	Hymexazol	Sankyo, Cleary	Drench	1 oz per acre 2 oz per acre	1 1
Metconazole	Caramba, Quash, V-10116	Tourney	BASF, Valent	Drench	3 fl oz per 100 gal 5.7 fl oz per 100 gal	1 8
				Foliar	5.7 fl oz per 100 gal	2
				Drench	1 oz per acre	1
				Foliar	1 oz per acre	1
Prothioconazole	Proline	Proline	Bayer	Drench	16 oz per 100 gal	3
				In-furrow	0.77 pt per 1000 ft	1
				Drench	10 fl oz per 100 gal	8
				Drip	10 fl oz per 100 gal	2
Pyraclostrobin	Cabrio, Headline	Insignia	BASF	Foliar	10 fl oz per acre	1
				Drench	5.3 fl oz per 100 gal 51.1 fl oz per 100 gal	3
				In-furrow	0.8 pt per 1000 ft	1
Thiophanate methyl	Topsin	3336	UPI, Cleary	Drench	9.5 fl oz per 100 gal 76.7 fl oz per 100 gal	3
				Drip		
				Foliar		
Tiadinil	Tiadinil		Nichino	Drench		
Trifloxystrobin	Flint, Gem	Compass	Bayer	In-furrow		
Triticonazole	Charter	Trinity	BASF	Drench		



## Results

### **Comparative Efficacy on *Fusarium avenaceum***

In 2001, McGovern conducted a trial to determine efficacy of several fungicides for control of *Fusarium* crown and stem rot (*F. avenaceum*) on lisianthus (*Eustoma grandiflorum*). All products were applied as a soil drench 24-hr prior to inoculation, and reapplied biweekly, except for Medallion which was reapplied after 1 month. Heritage, 3336 WP, Banner Maxx and a reduced-rate combination of Heritage and Medallion significantly reduced a severe *Fusarium* crown and stem rot incidence and plant mortality, with Heritage and Heritage + Medallion being the most effective (Table 3). Medallion and Fluazinam were less effective, and Systhane was ineffective. No phytotoxicity was observed from any treatment.

**Table 3. \* Efficacy on *Fusarium* Crown and Stem Rot (*Fusarium avenaceum*) on Lisianthus (*Eustoma grandiflorum*) ‘Maurine Blue’, McGovern, FL, 2001.**

Treatment	Rate Per 100 Gal	Final Disease Incidence (%) <sup>x</sup>	Final Mortality (%)	AUDPC <sup>y</sup>	AUMPC <sup>z</sup>
3336 50WP (thiophanate methyl)	16 oz	58.5 b	29.2 bc	522 cd	96 a
Banner Maxx 1.3MEC (propiconazole)	5 fl oz	66.0 b	33.5 b	1972 ab	965 a
Fluazinam 4.17F (fluazinam)	6.4 fl oz	70.8 ab	37.5 b	1425 bc	728 a
Heritage 50WG (azoxystrobin)	8.7 oz	20.8 c	0.0 c	314 d	0 a
Heritage + Medallion	4.35 + 1 oz	21.0 c	0.0 c	244 d	0 a
Medallion 50WP (fludioxonil)	2 oz	66.5 ab	29.2 bc	921 cd	2139 a
Systhane 40WSP (myclobutanil)	4 oz	91.8 ab	54.5 a	1978 a	1002 a
Untreated inoculated	-	100 a	87.5 a	2844 a	1457 a

\* Not an IR-4 Experiment: F&N Tests Vol 57: OT16.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.01).

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> AUMPC = area under the mortality progress curve.

### **Comparative Efficacy on *Fusarium commune***

In 2010 and 2011, Chastagner conducted three greenhouse trials to test the efficacy of several fungicides applied mainly as drench for control of damping off and root rot caused by *F. commune* (Isolates 34, 39, 53, 101, MBL12015) on Douglas fir (*Pseudotsuga menziesii*). In 2010, all products were applied 5 days after disease inoculation of potting mix (15 Dec), except BW240 which was applied on 12 Dec. Plants were seeded on 22 Dec. Treatments were applied one to three times on 1 to 4-week intervals for various products. In 2011, all treatments, with the exception of BW240, foliar Pageant, and the initial application of Acibenzolar were applied as drenches. On November 3, 2011 all treatments were planted with 10 seeds per pot. The initial application of Acibenzolar was applied directly to the seeds at this time. This was done by soaking the seeds in the Acibenzolar solution for 10 minutes prior to planting. The initial foliar application of Pageant was applied on November 21, 2011, which was 4-7 days after germination. Data on symptom development was collected once per week for 4 weeks beginning 14-15 days after seeding.

Notes on symptoms were taken and the number of “healthy” seedlings was recorded. Symptoms included damping off, which occurred shortly after emergence of the cotyledon, to root rot which killed the seedlings during the experiment. Disease pressure was high in all trials. In 2010, only the drench applications of Tourney at 1 or 2 oz/100 gallons had significantly higher numbers of healthy seedlings per pot (Table 4). Promising activity was observed for Compass, Pageant and Trinity as they were comparable to the non-inoculated check. In 2011, the drench applications of Torque, Pageant, 3336, CG100, Insignia at the high rate and Tourney had significantly higher numbers of healthy seedlings (Table 5). BW240 was ineffective (Table 6). No phytotoxicity was observed from any treatment except SP2169 in which some plants appeared to be stunted with malformed needles.

**Table 4. Efficacy on Damping-off and Root Rot Caused by *Fusarium commune* on Douglas Fir (*Pseudotsuga menziesii*), Chastagner WA, 2010.**

Treatment	Rate Per 100 Gal	Application Dates <sup>z</sup>	Application Interval	No. of Healthy Seedlings <sup>x,y</sup>
Acibenzolar	0.125 oz	2, 4	21 Days	2.2 b
	0.25 oz	2, 4	21 Days	2.4 b
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	6 oz	1	1 application	1.6 b
CG100 (caprylic acid)	0.6 pt	2, 5	28 Days	3.0 b
Compass (trifloxystrobin)	0.5 oz	2, 5	28 Days	4.0 ab
Heritage (azoxystrobin)	1.8 oz	2, 5	28 Days	4.0 ab
	4 oz	2, 5	28 Days	2.2 b
Hurricane (fludioxonil+mefenoxam)	0.75 oz	2	1 application	3.0 b
Medallion (fludioxonil)	2 oz	2, 4	21 Days	2.6 b
Pageant 38WG (boscalid+pyraclostrobin)	12 oz	2, 3, 5	14 Days	4.4 ab
Palladium (cyprodinil+fludioxonil)	6 oz	2, 3, 5	14 Days	1.8 b
Tourney (metconazole)	1 oz	2, 4	21 Days	8.2 a
	2 oz	2, 4	21 Days	8.2 a
Trinity (triticonazole)	6 fl oz	2, 3, 5	14 Days	5.2 ab
	8 fl oz	2, 3, 5	14 Days	5.0 ab
Untreated non-inoculated	-	-	-	5.7 ab
Untreated inoculated	-	-	-	1.9 b

<sup>x</sup> Means followed by the same letter do not differ significantly based on Tukey's HSD Test, (P=0.001).

<sup>y</sup> Data collected 35 days after seeding (January 26, 2011).

<sup>z</sup> Dates: 1 = 12/12/10, 2 = 12/20/10, 3 = 1/3/11, 4 = 1/10/11, 5 = 1/18/11.

**Table 5. Efficacy on Damping-off and Root Rot Caused by *Fusarium commune* on Douglas Fir (*Pseudotsuga menziesii*), Chastagner WA, 2011, Trial 1.**

Treatment	Rate Per 100 Gal	Application Method	App. Dates <sup>y</sup>	App. Interval	Emergence Out of 10 Seeds <sup>x</sup>	No. of Healthy Seedlings	No. of Plants W/ Phyto Symptoms
3336 F (thiophanate-methyl)	16 fl oz	Drench	1, 2, 5	14 Days	8.8 abc	7.0 abc	0.0 b
Acibenzolar-s-methyl	0.25 oz	Seed soak & Drench	2, 4	21 Days	6.2 bc	0.2 f	0.0 b
CG100 (caprylic acid)	0.6 pints	Drench	1, 5	28 Days	8.6 abc	5.0 bcd	0.0 b
Compass 50WDG (trifloxystrobin)	0.5 oz	Drench	1, 4	21 Days	8.8 abc	4.0cde	0.0 b
Heritage 50WG (azoxystrobin)	1.8 oz	Drench	1, 5	28 Days	7.4 abc	1.6 def	0.0 b
Hurricane (fludioxonil + mefenoxam)	0.75 oz	Drench	1	1 application	7.4 abc	1.6 def	0.0 b
Insignia SC (pyraclostrobin)	3.1 fl oz	Drench	1, 3, 5	14 Days	8.4 abc	2.4 def	0.0 b
	6.1 fl oz	Drench	1, 3, 5	14 Days	8.8 abc	4.8 bcd	0.0 b
Medallion (fludioxonil)	2 oz	Drench	1, 4	21 Days	7.8 abc	2.0 def	0.0 b
Pageant 38WG (boscalid+pyraclostrobin)	8 oz	Foliar	4	14 Days	7.2 abc	0.6 f	0.0 b
	12 oz	Drench	1, 3, 5	14 Days	9.4 a	7.2 abc	0.0 b
Palladium 62.5WG (cyprodinil + fludioxonil)	4 oz	Drench	1, 3, 5	14 Days	8.2 abc	2.4 def	0.0 b
SP2169	12.3 fl oz	Drench	1, 3, 5	14 Days	9.0 ab	4.4 cde	4.2 a
Torque (tebuconazole)	6 fl oz	Drench	1, 3, 5	14 Days	9.2 a	8.2 ab	0.0 b
Tourney 50 WDG (metconazole)	1 oz	Drench	1, 4	21 Days	8.8 abc	7.4 abc	0.0 b
	2 oz	Drench	1, 4	21 Days	9.8 a	9.0 a	0.0 b
Trinity 2 SC (triticonazole)	6 oz	Drench	1, 3, 5	14 Days	9.0 ab	2.4 def	0.0 b
	8 oz	Drench	1, 3, 5	14 Days	9.4 a	3.2 def	0.6 b
Untreated non-inoculated	-	-	-	-	8.8 abc	8.8 a	0.0 b
Untreated inoculated	-	-	-	-	6.0 c	1.2 ef	0.0 b

<sup>x</sup> Means followed by the same letter do not differ significantly based on Tukey's HSD Test, (P=0.001).

<sup>y</sup> Dates: 1 = 11/1/11, 2 = 11/3/11, 3 = 11/16/11, 4 = 11/21/11, 5 = 11/29/11.

Emergence data collected 28 days after seeding; no. of healthy seedlings and phytotoxicity collected 35 days after seeding (11/3/11).

**Table 6. Efficacy on Damping-off and Root Rot Caused by *Fusarium commune* on Douglas Fir (*Pseudotsuga menziesii*), Chastagner WA, 2011, Trial 2.**

Treatment	Rate Per 100 Gal	Application Method <sup>y</sup>	Emergence Out of 10 Seeds <sup>x</sup>	No. of Healthy Seedlings
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	8 oz	Soil incorp. and seed trt	0.0 b	0.0 b
Untreated non-inoculated	-	-	8.6 a	7.4 a
Untreated inoculated	-	-	7.6 a	0.4 b

<sup>x</sup> Means followed by the same letter do not differ significantly based on Tukey's HSD Test, (P=0.001).

<sup>y</sup> BW240 applied to soil 3 days before inoculation and to seeds before seeding 5 days after inoculation.

Emergence data collected 28 days after seeding; no. of healthy seedlings and phytotoxicity collected 35 days after seeding (11/3/11).

In 2012, Chastagner conducted two greenhouse trials to test the efficacy of several fungicides applied as pre-plant soil incorporation for control of damping off and root rot caused by *F. commune* (Isolate MBL12015) on Douglas fir and a natural infection on bulbous iris (*Iris x hollandica*). For the Douglas fir experiment, *Fusarium* inoculation to soil from a commercial grower's nursery field that had been routinely used for Douglas fir production was done prior to fungicide applications on November 30, except for the Rootshield Plus treatment which was applied to the soil on Nov 14. Ten seeds were planted for each pot on Dec 3. Based on isolations from symptomatic seedlings, the only treatment that did not yield any *Fusarium* was the field soil that was autoclaved. This suggests that the soil acquired from a commercial grower's nursery field was already infested with this pathogen. It is important to note that the steaming treatment used in this experiment did not eliminate the pathogen from the soil. The number of germinated seed, healthy seedlings and the height of healthy seedlings was highly variable from pot to pot within treatments during this test. Consequently, even though the number of germinated seeds and the height of the seedlings ranged from 4.3 to 8.0 and 2.4 to 6.0, respectively, there were no significant differences with these variables between any of the treatments (Table 7). The number of healthy seedlings ranged from 0.5 to 6.3. While there was still variability from pot to pot within treatments, the Flagship plus Heritage had a significantly higher number of healthy seedlings than the inoculated field soil, which is the appropriate check for this treatment. None the other fungicide treatments had any significant effect on the number of healthy seedlings.

For the bulbous iris experiment, fungicides were applied and incorporated on Aug 16 to soil that was acquired from a commercial grower's greenhouse that had been routinely used for iris production, except for Rootshield Plus which was applied and mixed on Aug 18. There were two checks included in the treatments. These were greenhouse soil that received treatment and greenhouse soil that was steamed for 2 hours at 200F to kill off inoculum in the soil. Bulbs were planted on August 21 (Reps 1 and 2) and August 22 (Reps 3, 4, 5). Disease pressure in this trial was high. Treatments had no effect on the number of iris plants that emerged (Table 8). The number of marketable flowers per crate ranged from 32.4 to 12.2 for the bulbs planted in the steam pasteurized and non-pasteurized soils, respectively. Bulbs planted in soil treated with the high rate of Trinity, Flagship 0.22 G + Heritage, and the high rate of Tourney had significantly higher numbers of marketable flowers, flowers >18" tall, and significantly fewer culls than the bulbs planted in the non-pasteurized soil. The combinations of Flagship 25WG + Heritage and Acibenzolar + Heritage also significantly increased the number of flowers >18" tall. Flagship 0.22 G alone significantly increased the total number of marketable flowers and the number of flowers >18" tall. *Fusarium* was isolated from 88% of the cull flower bulbs in the non-pasteurized soil. *Fusarium* was also isolated from 22% of the cull flower bulbs in the steam pasteurized soil, which suggests that the pathogen was present on some of the bulbs at the time of planting. This may have reduced the efficacy of some fungicide treatments.

**Table 7. Efficacy on Damping-off and Root Rot Caused by *Fusarium commune* on Douglas Fir (*Pseudotsuga menziesii*), Chastagner WA, 2012.**

Treatment	Rate/1000ft <sup>2</sup>	No. Germinated <sup>x</sup> 1/2/13	Healthy Seedlings 1/29/13	
			No. of Plants	Avg Ht (cm)
Acibenzolar + Heritage 50WDG	0.08g + 1oz	6.8 a	3.5 a-d	4.7 a
CG 100 (caprylic acid)	26.0 ml	6.5 a	2.8 bcd	3.7 a
Flagship 0.22G (thiamethoxam)	44 oz	6.3 a	4.0 abc	4.7 a
Flagship 0.22G +Heritage 50WDG	44 oz + 0.4oz	4.8 a	2.0 cd	4.5 a
Flagship 25WG (thiamethoxam)	0.55 oz	7.8 a	2.0 cd	4.9 a
Flagship 25WG + Heritage 50WDG	0.4 oz +0.4 oz	8.0 a	5.5 ab	4.2 a
Heritage 50WDG (azoxystrobin)	0.4 oz	5.8 a	3.5 a-d	5.0 a
Palladium 62.5WG (cyprodinil + fludioxonil)	1 oz	4.3 a	0.5 d	2.4 a
Rootshield Plus WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	10 oz	4.0 a	2.5 bcd	5.7 a
SP2169	8.35 ml	6.3 a	1.3 cd	3.2 a
Torque 3.6SC (tebuconazole)	20.4 ml	4.5 a	1.0 cd	3.9 a
Tourney 50WDG (metconazole)	2.8 g	5.8 a	3.5 a-d	3.8 a
	5.6 g	5.3 a	2.5 bcd	4.2 a
Trinity 2SC (triticonazole)	5.43 ml	5.8 a	3.0 a-d	4.3 a
	8.14 ml	5.8 a	2.0 cd	3.3 a
Autoclaved field soil	-	7.5 a	6.3 a	6.0 a
Steamed field soil	-	6.8 a	4.3 abc	5.5 a
Non-inoculated field soil	-	7.0 a	3.3 a-d	3.9 a
Inoculated field soil	-	7.3 a	1.8 cd	4.1 a

<sup>x</sup> Means followed by the same letter do not differ significantly based on Tukey's HSD Test, (P<0.05). Data collected 29 days after seeding (December 4, 2012).

**Table 8. Efficacy on Damping-off and Root Rot Caused by *Fusarium commune* on Bulbous Iris (*Iris x hollandica*) ‘Blue Diamond’, Chastagner WA, 2012.**

Treatment	Rate/1000ft <sup>2</sup>	No. Emerged <sup>x</sup> 9/5/12	No. of Flowers Per Crate		
			Marketable	18 Inch	Cull
Acibenzolar + Heritage 50WDG	0.08g + 1oz	44.4 a	22.4 a-d	21.6 abc	22.2 abc
CG 100 (caprylic acid)	26.0 ml	44.8 a	19.2 bcd	18.4 bcd	24.8 abc
Flagship 0.22G (thiamethoxam)	44 oz	45.0 a	23.0 abc	21.8 abc	22.0 abc
Flagship 0.22G +Heritage 50WDG	44 oz + 0.4oz	44.6 a	24.2 abc	23.4 ab	21.6 bc
Flagship 25WG	0.55 oz	44.4 a	17.6 bcd	17.4 bcd	26.8 abc
Flagship 25WG+Heritage 50WDG	0.4 oz +0.4 oz	44.8 a	22.6 a-d	21.8 abc	22.2 abc
Heritage 50WDG (azoxystrobin)	0.4 oz	44.6 a	19.0 bcd	18.6 bcd	25.0 abc
Palladium 62.5WG (cyprodinil+fludioxonil)	1 oz	44.4 a	21.2 bcd	20.8 bcd	22.8 abc
Rootshield Plus WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	10 oz	44.8 a	13.6 bcd	12.8 cd	31.4 ab
SP2169	8.35 ml	44.0 a	21.0 bcd	20.8 bcd	22.0 abc
Torque 3.6SC (tebuconazole)	20.4 ml	44.4 a	21.8 a-d	20.6 bcd	22.2 abc
Tourney 50WDG (metconazole)	2.8 g	44.6 a	21.4 bcd	20.8 bcd	22.4 abc
	5.6 g	44.4 a	23.8 abc	23.2 abc	20.4 cd
Trinity 2SC (triticonazole)	5.43 ml	44.8 a	21.2 bcd	21.0 bcd	23.0 abc
	8.14 ml	44.0 a	24.5 ab	24.0 ab	21.5 cd
Steam pasteurized soil	-	43.8 a	32.4 a	32.0 a	10.8 d
Non-pasteurized soil	-	45.0 a	12.2 d	11.0 d	32.2 a

<sup>x</sup> Means followed by the same letter do not differ significantly based on Tukey's HSD Test, (P<0.05). Data collected 14 days after planting bulbs (August 21-22, 2012).

<sup>y</sup> Harvested between October 9, 2012 and November 9, 2012.

### Comparative Efficacy on *Fusarium oxysporum*

In 2001, McGovern conducted two trials to determine efficacy of several fungicides for control of *Fusarium* wilt (*F. oxysporum*) on lisianthus (*Eustoma grandiflorum*). Treatments were applied as a soil drench 24 hr prior to inoculation with *F. oxysporum*, and reapplied at 2-3 wk intervals. In the first trial, severe final disease incidence was significantly reduced by Medallion and Hurricane (Table 9). Final plant mortality was significantly reduced by all treatments, with Medallion, and Hurricane the most effective in increasing plant survival. Plant height was significantly increased by Medallion, Hurricane and Compass. In the second trial, Medallion, Terraguard, Heritage, Systhane and 3336 WP significantly reduced a severe disease incidence and plant mortality, with Medallion and Terraguard being the most effective (Table 10). Spectro 90 was ineffective. No phytotoxicity was observed from any treatment.

**Table 9. \* Efficacy on *Fusarium* Wilt (*Fusarium oxysporum*) on Lisianthus (*Eustoma grandiflorum*) ‘Maurine Blue’, McGovern, FL, 2001.**

Treatment	Rate Per 100 Gal	Final Disease Incidence (%) <sup>x</sup>	Final Mortality (%)	AUDPC <sup>y</sup>	AUMPC <sup>z</sup>	Plant Height (cm)
Compass 50WDG (trifloxystrobin)	1 oz	88.9 a	39.4 bc	1698.0 b	508.5 b	18.2 c
Heritage 50WG (azoxystrobin)	0.9 oz	97.2 a	25.0 c	2085.7 b	295.0 bc	11.6 de
Hurricane 48WP (fludioxonil + mefenoxam)	1.5 oz	41.7 b	0.0 d	263.3 c	0.0 c	27.1 b
Medallion 50WP (fludioxonil)	2 oz	36.1 b	0.0 d	144.0 c	0.0 c	34.3 a
Untreated inoculated	-	100 a	69.4 a	3394.2 a	1220.2 a	10.5 de

\* Not an IR-4 Experiment: F&N Tests Vol 57: OT17. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.01).

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> AUMPC = area under the mortality progress curve.

**Table 10. \* Efficacy on *Fusarium* Wilt (*Fusarium oxysporum*) on Lisianthus (*Eustoma grandiflorum*) ‘Maurine Blue’, McGovern, FL, 2001.**

Treatment	Rate Per 100 Gal	Final Disease Incidence (%) <sup>x</sup>	Final Mortality (%)	AUDPC <sup>y</sup>	AUMPC <sup>z</sup>
3336 50WP (thiophanate methyl)	16 oz	75.0 b	33.3 b	502 b	161 b
Heritage 50WG (azoxystrobin)	8.7 oz	44.4 c	25.0 bc	326 bc	102 b
Medallion 50WP (fludioxonil)	2 oz	8.3 d	5.5 c	82 c	62 b
Spectro 90WDG (chlorothalonil + thiophanate methyl)	12 oz	88.8 ab	72.2 a	1560 a	948 a
Systhane 40WSP (myclobutanil)	4 oz	43.8 c	27.8 b	408 bc	304 b
Terraguard 50W (triflumizole)	6 oz	13.9 d	5.5 c	104 c	54 b
Untreated inoculated	-	94.4 a	77.8 a	1456 a	1055 a

\* Not an IR-4 Experiment: F&N Tests Vol 57: OT18.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.01).

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> AUMPC = area under the mortality progress curve.

In 2010, Kirk conducted a greenhouse trial to test the efficacy of several fungicides for control of root rot caused by *F. oxysporum* on larkspur (*Delphinium sp.*). All treatments were applied as drench, except BW240 which was applied to plant roots as an immersion in solution for 30 seconds prior to transplanting. Initial treatments were applied 4 days after transplanting, except Acibenzolar applied 14 days before, CG100 applied at transplanting and Trinity applied 21 days after transplanting. Plants were

inoculated immediately after the application of fungicides on 20 Sep. *Fusarium* necrosis and root rot developed in the trial and about 150 days after transplanting, the inoculated check plants developed severe leaf necrosis and root necrosis. All treatments except Tourney significantly reduced the foliar and root necrosis in comparison to the non-treated inoculated control (Table 11). It is possible that plants treated with Tourney were excessively inoculated. The treatments with the greatest efficacy included the standards 3336, and Medallion, with Hurricane almost comparable; Acibenzolar at the lower rate also looked promising. All treatments caused transient leaf phytotoxicity.



**Table 11. Efficacy on Fusarium Root Rot (*Fusarium oxysporum*) on Larkspur (*Delphinium sp.*), Kirk, MI, 2010.**

Treatment	Rate Per 100 Gal	Application Dates <sup>u</sup>	Phytotoxicity <sup>x, t</sup> 74 DAP <sup>v</sup>	No. Leaves Per Plant 104 DAP	Plant Height (cm) 104 DAP	Leaf Necrosis <sup>w</sup> 148 DAP	RAUDPC <sup>y</sup> 148 DAP	Root Necrosis <sup>z</sup> 148 DAP
Acibenzolar	0.125 oz	A	1.1 ab	9.4 de	30.3 de	39.0 ef	9.2 e	3.9 e
	0.25 oz	A	1.4 a	9.0 def	35.2 cd	47.5 d	10.3 d	4.6 cd
3336 80WG (thiophanate methyl)	4 lb	D	0.9 bc	11.2 bc	42.5 bc	28.5 h	8.8 e	2.6 f
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	6 oz, 3 oz	B, H	1.1 ab	7.5 fgh	25.5 de	56.0 c	11.9 c	4.4 cde
CG100 20SC (caprylic acid)	0.8 pt	C	0.9 bc	8.7 d-g	23.7 e	55.8 c	12.1 c	5.5 b
Hurricane (fludioxonil + mefenoxam)	12 oz	D	1.4 a	12.6 ab	51.3 ab	35.1 fg	9.2 e	2.6 f
Medallion (fludioxonil)	2 oz	D	1.0 bc	13.6 a	54.4 a	31.3 gh	9.3 e	2.7 f
Pageant 38WG (boscalid+pyraclostrobin)	12 oz	D, E	0.5 d	7.3 gh	28.0 de	46.3 d	11.1 cd	4.2 de
Tourney (metconazole)	2 oz	D	1.0 bc	3.0 i	41.6 bc	78.5 a	14.5 b	7.6 a
Trinity (triticonazole)	6 fl oz	F, G	0.7 cd	10.0 cd	35.3 cd	43.8 de	10.6 d	5.0 bc
Untreated non-inoculated	-	-	0.0 e	8.0 efg	34.2 cd	16.4 i	2.4 f	2.3 f
Untreated inoculated	-	-	0.0 e	6.2 h	23.5 e	65.0 b	17.6 a	7.7 a

<sup>x</sup> Means followed by the same letter do not differ significantly based on Fisher's LSD (P=0.05).

<sup>t</sup> Phytotoxicity scale from 0 – 5; 0= no phytotoxicity; 1= ≈1mm of entire leaf margin yellow of at least one leaf; 2= 1-5% of entire leaf margin yellow of at least one leaf; 3= 1-5% of entire leaf margin yellow of all leaves; 4= 5-10% of entire leaf margin yellow of all leaves; 5= >10% of entire leaf margin yellow of all leaves.

<sup>w</sup> Leaf necrosis percentage over whole plant.

<sup>y</sup> RAUDPC, relative area under the disease progress curve calculated from day of appearance of initial symptoms.

<sup>z</sup> Root necrosis scale from 0 – 10; 0= no necrosis; 1= 0-5%; 2= 6-10%; 3= 11-15%; 4= 16-20%; 5= 20-30%; 6= 30-40%; 7= 40-50%; 8= 50-60%; 9= 60-75%; 10= 75-100% of root mass necrotic.

<sup>v</sup> Days after planting

<sup>u</sup> Application dates: A= 2 Sep (2 weeks prior to inoculation); B= 16 Sep (root dip before planting); C= 16 Sep (soil drench at planting); D= 16 Sep; E= 30 Sep; F= 7 Oct; G= 21 Oct; H= 25 Nov

In 2011, Chase conducted a greenhouse trial to test the efficacy of several fungicides applied as drench/spreng for control of Fusarium wilt caused by *F. oxysporum* on lisianthus (*Eustoma grandiflora*). Plugs were planted on 21 March and all treatments applied as drench/spreng on 28 March. Plants were inoculated on 11 April. Additional treatments were applied at different intervals for various products on 4, 18 April, 2, 9, 16 and 23 May. Three weeks after test initiation, stunting due to either Fusarium or phytotoxicity had become apparent. The worst damage was seen with both rates of Tourney which caused severe stunting (Table 12). Only plants treated with acibenzolar at the low rate were as tall as the noninoculated control. After three weeks, top grade was lowest for plants treated with Tourney. After about four weeks, severity of Fusarium wilt was low for all treatments except for the inoculated control, BW240 and Phyton 27 alone. After six weeks, only the higher rate of acibenzolar, SP2169 and Tourney at the 1 oz rate showed significantly lower disease severity.

**Table 12. Efficacy on Fusarium wilt (*F. oxysporum*) on Lisianthus (*Eustoma grandiflora*), Chase, CA, 2011.**

Treatment	Rate per 100 Gal	Applic. Interval (Days)	Height (cm) 5-4-11	Top Grade 5-4-11	Disease Severity <sup>y</sup>		
					5-9-11	5-16-11	5-23-11
Acibenzolar	0.125 oz	21	13.2 d	3.2 f	1.0 a	1.0 a	1.3 ab
	0.25 oz	21	10.9 bcd	3.0 def	1.0 a	1.1 a	1.0 a
3336 80WG (thiophanate methyl)	16 oz	14	9.8 bc	2.9 cdef	1.2 a	1.1 a	1.7 abc
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	6 oz, then 3 oz		8.0 bc	2.7 bcd	1.9 ab	2.4 d	2.2 abc
CG100 20SC (caprylic acid)	9.6 fl oz	14	9.4 bc	2.8 bcdef	1.2 a	1.5 abcd	2.4 abc
Palladium (cyprodinil + fludioxonil)	6 oz	14	10.6 bcd	2.9 cdef	1.1 a	1.2 ab	1.4 ab
Phyton 27 (copper sulfate pentahydrate)	25 oz	14	9.7 bc	2.8 bcde	2.1 b	2.4 cd	2.9 c
Phyton 27 + X3	25 oz + 1:500	14	10.1 bcd	3.0 cdef	1.1 a	1.7 abcd	1.9 ab
SP-2169	12.3 fl oz	14	8.3 bc	2.8 bcdef	1.0 a	1.0 a	1.0 a
Tourney (metconazole)	1 oz	14	3.9 a	2.5 ab	1.2 a	1.2 abc	1.0 a
	2 oz	14	3.4 a	2.3 a	1.1 a	1.3 abcd	1.3 ab
Trinity (triticonazole)	6 fl oz	14	8.2 bc	2.9 cdef	1.1 a	1.3 abcd	1.6 abc
	8 fl oz	14	7.4 b	2.7 bcd	1.0 a	1.0 a	1.3 ab
X3 (hydrogen peroxide, peroxyacetic & octanoic acids)	1:500	14	8.2 bc	2.6 bc	1.3 a	1.5 abcd	2.1 abc
Untreated noninoculated			11.5 cd	3.1 ef	1.0 a	1.0 a	1.0 a
Untreated inoculated			9.6 bc	2.9 cdef	1.6 ab	2.3 bcd	2.7 bc

<sup>x</sup> Means followed by the same letter do not differ significantly (P=0.05).

<sup>y</sup> Disease severity was recorded using the following scale: 1 – no disease, 2 – slight, 3 – moderate, 4 – severe to 5 – plant dead.

In 2012, Kirk conducted two field trials to determine the efficacy of several fungicides for control of stem based necrosis caused by *F. oxysporum* on corn flag (*Gladiolus sp.*). In the first trial, corms were dipped in the fungicide-water solutions for one minute a day before planting on June 26. The corms were not inoculated. In the second trial, plots were inoculated with a conidial/water suspension of *F. oxysporum* f.sp. *gladioli* before planting corms on June 27. All treatments were applied twice on 25 Jul and Aug 15. In the first trial, Fusarium symptoms developed in the trial but did not greatly affect emergence, with the untreated check having 100% emergence 58 days after planting and generally no differences in emergence among treatments (Table 13). Acibenzolar, the lowest rate of Acibenzolar + Azoxystrobin and Disarm provided significant but poor reduction of stem based necrosis. However, overall data on Fusarium related symptoms showed no product providing consistently positive efficacy. The reduction in emergence, height and flowering rate may have been indicative of a physiological effect of the products applied. In the second trial, a check was planted in a non-inoculated area of the trial but more severe symptoms developed on these plants than on those in the inoculated area (Table 14). No treatments were significantly different from the untreated checks in terms of plant stand, % necrosis, plant height, lodging or flowering rate. In general, the plants from the foliar trial were visibly shorter than those in the corm dip trial.

**Table 13. Efficacy on Stem Based Necrosis Caused By *Fusarium oxysporum* on Corn Flag (*Gladiolus sp.*) ‘Sun-Kissed’, Kirk, MI, 2012, Trial 1.**

Treatment	Rate Per 100 Gal	Plant Stand (%) <sup>x</sup>				RAUEPC <sup>y</sup> 58 DAP	Fusarium Related Symptoms			
		21 DAP <sup>u</sup>	28 DAP	58 DAP	89 DAP		% Necrosis <sup>w</sup> 59 DAP	Plant Height (in) 59 DAP	% Lodging <sup>y</sup> 69 DAP	% Flowering <sup>z</sup> 89 DAP
3336 WP (thiophanate methyl)	16 oz	7.1 cd	67.9 a-d	96.4 ab	90.1 a-d	52.0 ab	28.6 abc	20.7 bc	28.6 def	50.0 c
Acibenzolar 50WG	0.125 oz	3.6 d	28.6 e	92.9 b	82.1 d	54.8 ab	17.9 c	19.5 c	57.1 abc	57.1 bc
	0.25 oz	35.7 ab	71.4 a-d	100.0 a	100.0 a	53.8 ab	21.4 bc	22.0 bc	50.0 a-e	60.7 abc
Acibenzolar + Heritage (azoxystrobin)	0.25 oz + 0.9 oz	10.7 bcd	78.6 abc	100.0 a	100.0 a	57.7 ab	28.6 abc	23.9 ab	60.7 ab	53.6 c
	0.125 oz + 0.45 oz	3.6 d	60.7 a-e	92.9 b	92.9 abc	46.0 b	10.7 c	23.3 bc	71.4 a	75.0 abc
	0.5 oz + 1.8 oz	21.4 a-d	82.1 ab	100.0 a	96.4 ab	63.0 ab	21.4 bc	22.4 bc	46.4 a-e	67.9 abc
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	80 oz	3.6 d	46.4 cde	100.0 a	92.9 abc	59.4 ab	21.4 bc	21.5 bc	35.7 b-f	85.7 a
CG100 20SC (caprylic acid)	38.4 fl oz	35.7 ab	89.3 a	96.4 ab	92.9 abc	59.1 ab	28.6 abc	21.8 bc	17.9 f	82.1 ab
Disarm (fluoxastrobin)	4 fl oz	32.1 abc	78.6 abc	100.0 a	92.9 abc	60.3 ab	14.3 c	23.9 ab	25.0 ef	64.3 abc
Medallion (fludioxonil)	8 oz	7.1 cd	75.0 abc	100.0 a	89.3 bcd	54.3 ab	32.1 abc	22.1 bc	53.6 a-d	75.0 abc
Palladium (cyprodinil + fludioxonil)	6 fl oz	10.7 bcd	50.0 b-e	96.4 ab	82.1 d	48.5 b	25.0 bc	21.6 bc	35.7 b-f	71.4 abc
SP-2169	12.3 fl oz	39.3 a	89.3 a	96.4 ab	100.0 a	61.4 ab	21.4 bc	22.1 bc	53.6 a-d	64.3 abc
Tourney (metconazole)	2 fl oz	7.1 cd	39.3 de	100.0 a	96.4 ab	54.7 ab	50.0 a	21.9 bc	60.7 ab	85.7 a
	4 fl oz	14.3 a-d	85.7 a	96.4 ab	85.7 cd	54.8 ab	32.1 abc	21.2 bc	32.1 c-f	50.0 c
Untreated	-	35.7 ab	85.7 a	100.0 a	96.4 ab	66.3 a	42.9 ab	28.1 a	42.9 b-f	53.6 c

<sup>x</sup> Means followed by the same letter do not differ significantly based on Fisher’s LSD (P=0.05).

<sup>w</sup> Incidence of plants with visible stem base necrosis at least up to 0.5” above the soil line.

<sup>y</sup> Incidence of plants greater than 45° bending from the perpendicular.

<sup>z</sup> The percentage of plants with at least one inflorescence emerging from stem.

<sup>u</sup> Days after planting

<sup>v</sup> RAUDPC, relative area under the emergence progress curve.

**Table 14. Efficacy on Stem Based Necrosis Caused By *Fusarium oxysporum* on Corn Flag (*Gladiolus sp.*) ‘Sun-Kissed’, Kirk, MI, 2012, Trial 2.**

Treatment	Rate Per 100 Gal	Plant Stand (%) <sup>x</sup>		Fusarium Related Symptoms			
		58 DAP <sup>v</sup>	89 DAP	% Necrosis <sup>w</sup> 59 DAP	Plant Height (in) 59 DAP	% Lodging <sup>y</sup> 69 DAP	% Flowering <sup>z</sup> 89 DAP
3336 EC (thiophanate methyl)	16 fl oz	75.0 b	75.0 ab	21.4 a-d	14.0 abc	28.6 ab	39.3 a
Acibenzolar 50WG 0.125 oz	0.50 oz	75.0 b	71.4 b	14.3 cd	14.3 abc	35.7 ab	21.4 ab
	0.75 oz	89.3 ab	75.0 ab	46.4 a	13.5 abc	50.0 a	21.4 ab
Acibenzolar + Heritage (azoxystrobin)	0.5 oz + 1.8 oz	82.1 ab	82.1 ab	42.9 ab	13.5 abc	39.3 ab	25.0 ab
	0.75 oz + 2.7 oz	92.9 ab	89.3 ab	46.4 a	12.5 bc	35.7 ab	21.4 ab
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	8 oz	89.3 ab	85.7 ab	32.1 a-d	15.2 abc	46.4 ab	14.3 b
CG100 20SC (caprylic acid)	38.4 fl oz	96.4 a	82.1 ab	39.3 abc	13.6 abc	32.1 ab	32.1 ab
Disarm (fluoxastrobin)	4 fl oz	92.9 ab	89.3 ab	7.1 d	14.0 abc	35.7 ab	28.6 ab
Medallion (fludioxonil)	4 oz	85.7 ab	75.0 ab	17.9 bcd	16.7 ab	35.7 ab	21.4 ab
Palladium (cyprodinil + fludioxonil)	6 fl oz	96.4 a	92.9 ab	25.0 a-d	11.9 c	28.6 ab	17.9 ab
SP-2169	12.3 fl oz	96.4 a	96.4 a	25.0 a-d	17.0 a	25.0 b	35.7 ab
Tourney (metconazole)	1 fl oz	89.3 ab	89.3 ab	35.7 abc	13.4 abc	25.0 b	14.3 b
	2 fl oz	85.7 ab	82.1 ab	7.1 d	14.2 abc	35.7 ab	21.4 ab
Trinity (triticonazole)	8 fl oz	96.4 a	92.9 ab	35.7 abc	14.1 abc	39.3 ab	39.3 a
	12 fl oz	85.7 ab	82.1 ab	25.0 a-d	17.6 a	25.0 b	25.0 ab
Untreated non-inoculated	-	96.4 a	78.6 ab	39.3 abc	14.0 abc	32.1 ab	21.4 ab
Untreated inoculated	-	89.3 ab	82.1 ab	17.9 bcd	14.4 abc	25.0 b	25.0 ab

<sup>x</sup> Means followed by the same letter do not differ significantly based on Fisher’s LSD (P=0.05).

<sup>w</sup> Incidence of plants with visible stem base necrosis at least up to 0.5” above the soil line.

<sup>y</sup> Incidence of plants greater than 45° bending from the perpendicular.

<sup>z</sup> The percentage of plants with at least one inflorescence emerging from stem.

<sup>v</sup> Days after planting

From 2007 to 2008, five greenhouse studies were conducted to determine efficacy of several fungicides for control of Fusarium wilt (*F. oxysporum* f. sp. *niveum*) on watermelon (*Citrullus lanatus*). Treatments were applied as soil drench 72 hr prior to soil drench inoculation with *F. oxysporum*. In two trials conducted by Egel in 2007, Topsin provided the best control of Fusarium wilt, with Actigard and Tachigaren at the higher rate less effective (Table 15). Cannonball and Tachigaren at the lower rate provided significant control in one of two trials. No phytotoxicity was observed from treatments except Actigard which reduced seedling growth. In a 2008 experiment by Egel, Topsin, V-10116, Proline and Tiadinil provided 100 % control of a severe Fusarium wilt pressure, similar to uninoculated control (Table 16). Quadris was essentially as effective as these products, Charter and Cabrio were less effective and Actigard and Cannonball virtually ineffective. No phytotoxicity was observed from any treatment in this trial.

**Table 15. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Egel, IN, 2007.**

Treatment	Rate Per Acre <sup>w</sup>	Test 1 Feb 2007		Test 2 March 2007	
		AUDPC <sup>x,y</sup>	Plant Size (cm <sup>2</sup> ) <sup>z</sup> 3/12/07	AUDPC	Plant Size (cm <sup>2</sup> ) 4/6/07
Actigard 50WG (acibenzolar)	0.33 oz	220.90 c	118.8 c	140.42 c	350.4 c
Cannonball 50WP (fludioxonil)	0.5 lb	607.62 a	310.2 a	353.51 b	756.8 a
Tachigaren 30L (hymexazol)	3 ml/sq m	365.37 b	311.7 a	563.76 a	712.4 a
Tachigaren 30L (hymexazol)	6 ml/sq m	252.13 c	307.5 a	348.22 b	653.8 ab
Topsin 4.5FL (thiophanate methyl)	10 fl oz	15.35 d	213.1 b	0.00 d	637.3 ab
Untreated non-inoculated	-	0.00 d	237.2 ab	0.00 d	501.8 bc
Untreated inoculated	-	679.89 a	-	531.10 a	-

\* Not an IR-4 Experiment: IR-4 Food Crops Website (online).

<sup>x</sup> Means followed by same letter do not differ significantly (P=0.05, LSD).

<sup>w</sup> Rate per acre was calculated on the assumption of 1,400 plants per acre.

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> The height and the maximum width of the leaves were measured in cm and the result multiplied to give an area.

**Table 16. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Egel, IN, 2008.**

Treatment	Rate Per 100 Gal	AUDPC <sup>x,y</sup>	Incidence <sup>z</sup>
Actigard 50WG (acibenzolar)	0.75 oz	32 b	0.74 ab
Cabrio EG (pyraclostrobin)	16 oz	91 b	0.25 de
Cannonball 50WP (fludioxonil)	16 oz	659 a	0.66 bc
Charter 25FS (triticonazole)	76.7 fl oz	45 b	0.16 de
Proline 480SC (prothioconazole)	5.7 fl oz	0 b	0 e
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	6 b	0.08 e
Tiadinil (tiadinil)	51.1 fl oz	0 b	0 e
Topsin 4.5FL (thiophanate methyl)	10 fl oz	0 b	0 e
V-10116 (metconazole)	8 oz	0 b	0 e
Untreated non-inoculated	-	0 b	0 e
Untreated inoculated	-	646 a	1.00 a

\* Not an IR-4 Experiment: IR-4 Food Crops Website (online). Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly (P=0.01, LSD).

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> 1.00 = 100 % disease incidence.

In a greenhouse trial conducted by Langston in 2008, Actigard, V-10116, Quadris, Proline and Topsin significantly reduced a very high Fusarium wilt pressure (Table 17). Significant stunting and phytotoxicity were observed in all treatments except Proline, with V-10116 treatments showing the most severe phytotoxicity. These data indicate that Proline, Actigard, Quadris and Topsin may have potential in watermelon production as at-planting applications for suppressing losses to Fusarium wilt. A greenhouse study conducted by Everts in 2008 showed that Quadris, Actigard at either high and low rate, Topsin, Proline and Metconazole were highly effective in reducing a very high Fusarium wilt pressure (Table 18). No phytotoxicity was observed for all treatments except Actigard and Metconazole. Actigard caused necrotic lesions, leaf yellowing, and slow plant growth although the symptoms were less with the low rate. Metconazole caused stunted plants with dark green leaves.

**Table 17. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Langston, GA, 2008.**

Treatment	Rate Per 100 Gal	Disease Severity <sup>x,y</sup> 5/12/08	AUDPC <sup>z</sup>
Actigard 50WG (acibenzolar)	0.75 oz	3.2 c	34.2 e
Cabrio 20EG (pyraclostrobin)	16 oz	9.7 a	100.3 a-d
Cannonball 50WP (fludioxonil)	16 oz	9.8 a	105.2 a-c
Proline 4SC (prothioconazole)	5.7 fl oz	5.9 b	50.3 e
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	5.5 b	46.6 e
Tiadinil 30% SC (tiadinil)	51.1 fl oz	9.9 a	112.8 a
Topsin 4.5F (thiophanate methyl)	10 fl oz	8.5 a	79.5 d
Trinity 1.69SC (triticonazole)	9.5 fl oz	9.9 a	109.0 a-c
V-10116 50WG (metconazole)	4 oz	5.2 bc	40.8 e
Untreated inoculated	-	9.8 a	108.3 a-c

\* Not an IR-4 Experiment: PDM Reports Vol 3: V155. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

<sup>y</sup> 0-10 scale where 0= no foliar wilt symptoms, 5.0=50% of the foliage wilted, and 10.0= a dead plant.

<sup>z</sup>. AUDPC = area under the disease progress curve calculated from severity ratings taken on May 1, 5, 8, 12 and 15.

**Table 18. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Sugar Baby’, Everts, MD, 2008.**

Treatment	Rate Per 100 Gal	Wilt Incidence (%) <sup>x</sup>	Final Wilt Severity	AUWSPC (%.days) <sup>y</sup>
Actigard 50WG (acibenzolar)	0.25 oz	8 bc	1 cd	6 e
Actigard	0.75 oz	0 c	0 d	0 e
Cabrio 20EG (pyraclostrobin)	16 oz	58 a	68 ab	635 cd
Cannonball 50WP (fludioxonil)	16 oz	75 a	62 ab	707 bcd
Metconazole 50WG (metconazole)	4 oz	17 b	6 cd	16 e
Proline 480SC (prothioconazole)	5.7 fl oz	17 b	2 cd	12 e
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	0 c	0 d	0 e
Tiadinil 30% SC (tiadinil)	5.3 fl oz	100 a	67 ab	977 bc
Topsin 4.5FL (thiophanate methyl)	10 fl oz	17 b	1 cd	1 e
Trinity 1.69SC (triticonazole)	9.5 fl oz	83 a	49 b	438 d
Untreated inoculated	-	100 a	99 a	1510 a

\* Not an IR-4 Experiment: PDM Reports Vol 3: V092. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

<sup>y</sup>. AUWSPC = area under wilt severity progress curve.

Four field experiments were conducted from 2008 to 2009 to determine efficacy of several fungicides for control of natural populations of Fusarium wilt (*F. oxysporum* f. sp. *niveum*) on watermelon (*Citrullus lanatus*). Everts conducted two trials where treatments were drenched to the soil around each plant immediately after transplanting. In the first trial planted 10 June, Actigard, Proline and Topsin significantly reduced wilt at 2½ weeks after transplanting (Table 19). The same three treatments that reduced wilt had the highest numerical vine length, although there were no significant differences among treatments. No phytotoxicity was observed for all treatments except Metconazole causing stunting and dark green leaves early in the season. In a second trial planted 3 June, all treatments, except Quadris, were effective in reducing wilt incidence at both 4 and 5 weeks after transplanting (Table 20). Watermelons treated with Actigard, Proline, Topsin and Quadris had longer vines than the untreated control. No phytotoxicity was observed for any treatment except Metconazole causing a significant

reduction in growth of plants, and dark green leaves during the early growing season. In 2009, Everts determined the efficacy of Actigard, Proline and Topsin applied through drip irrigation immediately after transplanting on 16 June, and again at 2 and 4 weeks after transplanting. Proline was the only product that significantly reduced a severe Fusarium wilt incidence where untreated plots had 78 % of plants with wilt symptoms by 1 Aug. (Table 21). Plot vigor and yield were numerically highest in plots where Proline was applied alone or in combination with other fungicides, although not significantly higher than in some treatments. No phytotoxicity was observed in any treatment.

**Table 19. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Sugar Baby’, Everts, DE, 2008.**

Treatment	Rate Per 100 Gal	Wilt Incidence (%) <sup>x</sup> at		Vine Length (in) 7/11/08	Plant Vigor (%) 7/22/08	Marketable Fruit	
		2 ½ wk	4 ½ wk			T/A	No./A
Actigard 50WG (acibenzolar)	0.25 oz	5 b	5	43	94	9.8	3449
Metconazole 50WG (metconazole)	4 oz	18 ab	15	30	76	7.0	2723
Proline 480SC (prothioconazole)	5.7 fl oz	5 b	10	44	92	7.5	3086
Quadris F (azoxystrobin)	15.4 fl oz	18 ab	20	33	72	6.6	2783
Topsin 4.5FL (thiophanate methyl)	10 fl oz	8 b	8	37	90	7.4	2602
Untreated	-	25 a	23	34	79	6.5	3086
<i>P</i> Values		0.0498	0.2893	0.0551	0.0516	0.4536	0.3631

\* Not an IR-4 Experiment: PDM Reports Vol 3: V097. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).



**Table 20. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Sugar Baby’, Everts, MD, 2008.**

Treatment	Rate Per 100 Gal	Wilt Incidence (%) <sup>x</sup> at			AUWIPC (% days) <sup>y</sup>	Wilt Severity (%) <sup>z</sup>	Vine Length (in) 6/27/08	Marketable Fruit 7/29/08	
		4 wk	5 wk	6 wk				T/A	No./A
Actigard 50WG (acibenzolar)	0.25 oz	0 b	2 b	25	734	61	72 ab	8.6	2133
Metconazole 50WG (metconazole)	4 oz	0 b	8 b	35	915	71	49 d	9.1	2042
Proline 480SC (prothioconazole)	5.7 fl oz	2 b	8 b	37	909	66	72 ab	8.7	2224
Quadris F (azoxystrobin)	15.4 fl oz	4 ab	15 ab	42	1057	60	70 ab	6.5	1679
Topsin 4.5FL (thiophanate methyl)	10 fl oz	2 b	6 b	33	875	63	73 a	8.5	2269
Untreated	-	8 a	29 a	58	1494	77	64 c	5.9	1588
<i>P</i> Values		0.0460	0.0212	0.2089	0.1267	0.2019	<0.0001	0.2956	0.4318

\* Not an IR-4 Experiment: PDM Reports Vol 3: V093. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

<sup>y</sup> The area under wilt incidence progress curve (AUWIPC) was calculated from wilt incidence data made at 3 through 8 weeks after transplanting.

<sup>z</sup> Final wilt severity was rated at fruit harvest using the Horsfall-Barratt scale.

**Table 21. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Sugar Baby’, Everts, MD, 2009.**

Treatment	Rate Per 100 Gal	Wilt Count Per Plot <sup>x</sup>		Plant Vigor (%) 8/7/09	Fruit Wt (Lb/Plot) 8/12/09
		7/17/09	8/1/09		
Actigard 50WG (acibenzolar)	0.25 oz	13 a	12.3 a	35.5 bcd	36.5 e
Actigard + Proline	0.25 oz + 5.7 fl oz	7.0 cd	8.5 b	38.0 bc	100.3 abc
Actigard + Topsin	0.25 oz + 10 fl oz	9.8 bc	12.3 a	26.0 cde	75.4 bcd
Proline 480SC (prothioconazole)	5.7 fl oz	5.0 d	6.3 b	45.0 b	105.7 a
Proline + Topsin	5.7 fl oz + 10 fl oz	7.3 cd	7.8 b	47.5 ab	102.8 ab
Topsin 4.5FL (thiophanate methyl)	10 fl oz	10.8 ab	14.0 a	19.0 e	73.4 cd
Untreated	-	10.0 abc	14.0 a	19.0 e	80.8 a-d

\* Not an IR-4 Experiment: PDM Reports Vol 4: V065. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

In 2009, Engel conducted a field trial to determine efficacy of Actigard, Proline and Topsin applied through trickle irrigation. All products were equally effective against Fusarium wilt (Table 22). No phytotoxicity was observed in any treatment.

**Table 22. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Imagination’, Egel, IN, 2009.**

Treatment <sup>w</sup>	Rate Per Acre	AUDPC <sup>x,y</sup>	Fruit Weight (Lb/A) <sup>z</sup>
Actigard 50WG (acibenzolar)	0.25 oz	257.4 c	32,125 a
Proline 480SC (prothioconazole)	5.7 fl oz	233.6 c	32,307 a
Topsin 4.5FL (thiophanate methyl)	10 fl oz	352.3 bc	31,581 a
Untreated Control	-	691.2 a	27,388 a

\* Not an IR-4 Experiment: IR-4 Food Crops Website (online). Not all products tested included in table.

<sup>w</sup> Applied 12, 26 May and 10 June.

<sup>x</sup> Means followed by same letter do not differ significantly (P=0.05,LSD).

<sup>y</sup> AUDPC = area under the disease incidence progress curve.

<sup>z</sup> Fruit was harvested 20, 27 July, 3 and 10 August.

In 2010, Langston conducted a field trial to determine efficacy of Actigard, Proline, Quadris, Topsin and V-10116 applied as drench at transplanting (4/7) followed by foliar spray 4 weeks later (5/5). All products significantly reduced Fusarium wilt incidence, with Actigard, Proline and V-10116 at the high rate providing superior control (Table 23). Significant phytotoxicity was observed with Actigard, Proline and V-10116, with V-10116 causing severe stunting. There was no difference in yields between treatments (data not shown).

**Table 23. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Langston, GA, 2010.**

Treatment <sup>w</sup>	Rate <sup>y</sup>	Plant Vigor <sup>z</sup> 4/21	% Wilt Incidence <sup>x</sup>	
			4/21	5/10
Actigard 50WG (acibenzolar)	0.75 oz	7.3 b	0.0 b	11.2 cd
Proline 4SC (prothioconazole)	5.7 fl oz	6.3 c	0.0 b	5.7 d
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	8.7 a	0.0 b	33.2 bc
Topsin 4.5F (thiophanate methyl)	10 fl oz	8.8 a	0.0 b	55.7 b
V-10116 50WG (metconazole)	1 oz	4.7 d	2.7 b	30.7 c
	2 oz	3.3 e	0.0 b	5.5 d
Untreated inoculated	-	8.7 a	33.3 a	94.3 a

\* Not an IR-4 Experiment: PDM Reports Vol 5: V156. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

% wilt incidence was rated by counting the number of plants in each plot that showed signs of wilting and dividing that number by the total number of plants in each plot x100.

<sup>w</sup> Products were applied on 4/7 as transplant drench; 5/5 as foliar spray.

<sup>y</sup> Rates are in per 100 gal as transplant drench and per acre as foliar spray.

<sup>z</sup> Plant Vigor was rated on 1-10 scale where 1= a dead or dying plant, 5 = moderately stunted plant and 10 = a healthy non-stunted plant.

In 2011, Langston conducted 3 field trials to determine the efficacy of Actigard, Proline, Quadris, and Serenade. In the first 2 trials, plots were inoculated with *Fusarium* before transplanting, then products were applied as drench at transplanting. In the first trial, Quadris and Serenade provided poor and no control of a severe Fusarium wilt incidence, resulting in plant vigor and yield that were not different from inoculated check (Table 24).

**Table 24. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Langston, GA, 2011, Trial 1.**

Treatment <sup>w</sup>	Rate Per 100 Gal	Plant Vigor <sup>y</sup> 4/1	% Wilt Incidence <sup>x</sup>		Yield (Lb/Plot)
			4/7	4/21	
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	5.8 a	58.3 c	78.3 a	28.8 a
Serenade ( <i>Bacillus subtilis</i> )	2 qt	6.7 a	95.0 a	96.7 a	10.8 a
	4 qt	6.7 a	96.7 a	95.0 a	3.5 a
Untreated inoculated	-	6.8 a	86.7 ab	95.0 a	14.5 a

\* Not an IR-4 Experiment: PDM Reports Vol 6: V138. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

% wilt incidence was rated by counting the number of plants in each plot that showed signs of wilting and dividing that number by the total number of plants in each plot x100.

<sup>w</sup> Products applied as plant drenches at transplanting on 3/17.

<sup>y</sup> Plant Vigor was rated on 1-10 scale where 1= a dead or dying plant, 5 = moderately stunted plant and 10 = a healthy non-stunted plant.

In the second trial, all products significantly reduced a severe Fusarium wilt incidence, with Actigard at the high rate providing the best control (Table 25). All treatments had significantly higher stand counts but did not significantly increase yield. In the third trial, products were applied as incorporated spray applications a week before transplanting or drench applied at transplanting in a field that had a history of severe Fusarium wilt. The disease progressed rapidly and became severe before the watermelons reached maturity, and no harvestable fruit was produced in any plots. All treatments significantly reduced plant vigor when applied as drench but not when applied as soil incorporated treatments (Table 26). All of the

drench treatments, and only the Proline soil incorporated treatment significantly reduced Fusarium wilt incidence. Only Proline and Actigard applied as drenches significantly reduced stand loss.

**Table 25. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Langston, GA, 2011, Trial 2.**

Treatment <sup>w</sup>	Rate Per 100 Gal	Plant Vigor <sup>y</sup> 4/1	% Wilt Incidence <sup>x</sup>		Stand Count 4/27	Yield (Lb/Plot)
			4/7	4/21		
Actigard 50WG (acibenzolar)	0.25 oz	4.5 a	38.3 c	86.7 a-c	1.3 de	24.7 a
	0.50 oz	4.3 a	21.7 cd	63.3 b-e	4.2 ab	36.5 a
	0.75 oz	4.5 a	11.7 d	41.6 e	4.3 ab	37.2 a
Proline 4SC (prothioconazole)	3.0 fl oz	4.7 a	30.0 cd	70.0 b-d	4.0 ab	29.8 a
	5.7 fl oz	5.0 a	36.7 cd	55.0 de	3.7 ab	40.3 a
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	5.0 a	36.7 cd	88.3 ab	1.8 cd	28.7 a
Untreated inoculated	-	5.3 a	100.0 a	100.0 a	0.0 e	0.0 a

\* Not an IR-4 Experiment: PDM Reports Vol 6: V154. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

% wilt incidence was rated by counting the number of plants in each plot that showed signs of wilting and dividing that number by the total number of plants in each plot x100.

<sup>w</sup> Products applied as plant drenches at transplanting on 3/17.

<sup>y</sup> Plant Vigor was rated on 1-10 scale where 1= a dead or dying plant, 5 = moderately stunted plant and 10 = a healthy non-stunted plant.

**Table 26. \* Efficacy on Fusarium Wilt (*Fusarium oxysporum* f. sp. *niveum*) on Watermelon (*Citrullus lanatus*) ‘Black Diamond’, Langston, GA, 2011, Trial 3.**

Treatment <sup>w</sup>	Rate <sup>y</sup>	Application Method	Plant Vigor <sup>z</sup> 3/28	% Wilt Incidence <sup>x</sup> 4/8	Stand Count 4/13
Actigard 50WG (acibenzolar)	0.50 oz	Drench	4.8 c	20.0 d	5.3 a
	0.25 oz	Incorp. spray	7.5 a	53.3 a-c	4.2 a-c
Proline 4SC (prothioconazole)	5.7 fl oz	Drench	6.0 b	33.3 cd	5.7 a
	5.7 fl oz	Incorp. spray	7.2 a	45.0 bc	4.7 ab
Quadris 2.08SC (azoxystrobin)	15.4 fl oz	Drench	5.8 bc	36.7 cd	4.5 ab
	15.4 fl oz	Incorp. spray	7.5 a	63.3 ab	2.7 c
Untreated inoculated	-		7.2 a	70.0 a	3.5 bc

\* Not an IR-4 Experiment: PDM Reports Vol 6: V139. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD (P=0.05).

% wilt incidence was rated by counting the number of plants in each plot that showed signs of wilting and dividing that number by the total number of plants in each plot x100.

<sup>w</sup> Incorporated spray applications applied on 3/8 and drench applications applied after transplanting on 3/14.

<sup>y</sup> Rates were on per acre for incorporated spray applications and on per 100 gal for drench applications.

<sup>z</sup> Plant Vigor was rated on 1-10 scale where 1= a dead or dying plant, 5 = moderately stunted plant and 10 = a healthy non-stunted plant.

### **Comparative Efficacy on *Fusarium solani***

In 2001, McGovern conducted a trial to determine efficacy of several fungicides for control of Fusarium tuber rot (*F. solani*) on caladium (*Caladium x hortulanum*). Tubers were soaked overnight (~16 hr) at ambient temperature in each of the fungicides on 16 May 2000, 3 days prior to planting. Tubers to be sprayed at planting with Hurricane initially were soaked overnight in water alone. Hurricane was applied at planting on 19 May 2000 with a hand-powered hydraulic sprayer and an approximate volume of the

diluted fungicide of 1500 gal/A. Plant emergence was significantly increased compared to the water control by preplant soak application of Spectro and Heritage and by Hurricane applied at planting (Table 27). Spectro significantly increased the number of marketable tubers per plot. Spectro, Heritage at 16 oz, or Hurricane applied as tuber soaks, and spray application of Hurricane at planting significantly increased total tuber weight per plot. Hurricane applied as a tuber soak or as a spray at planting was the only product that significantly reduced the severity of *Fusarium* tuber rot. No phytotoxicity was observed from any treatment.

**Table 27. \* Efficacy on Fusarium Tuber Rot (*Fusarium solani*) on Caladium (*Caladium x hortulanum*) ‘Florida Cardinal’, McGovern, FL, 2001.**

Treatment <sup>y</sup>	Rate Per 100 Gal	% Emergence <sup>x</sup> 6/29/00	No Marketable Tubers 4/9/01	Tuber Weight (Lb/Plot) 4/9/01	Disease Severity <sup>z</sup> 3/27/01
3336 50WP (thiophanate methyl)	24 oz	67.7 ab	71.8 e	8.0 d	8.8 abcd
Compass 50WDG (trifloxystrobin)	4 oz	72.5 bc	78.2 bcde	8.8 bcd	7.5 bcd
Heritage 50WG (azoxystrobin)	4 oz	63.6 a	76.0 cde	8.8 bcd	10.6 a
Heritage	8 oz	77.7 cde	82.2 abcde	8.6 bcd	9.0 abc
Heritage	16 oz	74.2 bcd	90.6 ab	9.6 ab	9.3 abc
Hurricane 48WP (fludioxonil + mefenoxam)	1.5 oz	76.0 cd	87.5 abcd	10.1 a	6.3 d
Hurricane spray at planting	1.5 oz	85.7 f	83.4 abcde	9.7 ab	7.0 cd
Spectro 90WDG (chlorothalonil +thiophanate methyl)	24 oz	83.3 ef	92.6 a	9.8 ab	9.5 abc
Untreated	-	71.0 bc	77.8 bcde	8.2 cd	9.6 ab

\* Not an IR-4 Experiment: F&N Tests Vol 57: OT05. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Fisher’s Protected LSD.

<sup>y</sup> All fungicides were applied as tuber soaks unless otherwise indicated.

<sup>z</sup> Tuber rot severity was measured using a 1-5 rating scale where 1 = 0%, 2 = 1-10%, 3 = 11-25%, 4 = 26-50%, and 5 = 51-100% internal discoloration.

In 2002, Kirk conducted a field trial to determine efficacy of BAS 500, Bravo Zn, Gem and Quadris for control of root rot caused by *Fusarium solani* f. sp. *phaseoli* on snap bean (*Phaseolus vulgaris*). All plots were inoculated and in-furrow applications of fungicides were made over the seed at planting. No treatments were significantly different from each other as measured by root rot index at the first harvest (32 DAP) but at the second harvest (72 DAP) all treatments had a significantly lower root rot index value than the untreated control (Table 28). No treatments were significantly different from each other in plant emergence and marketable yield. No phytotoxicity was observed from any treatment.

In 2011, Palmateer conducted a greenhouse trial to test the efficacy of several fungicides for control of stem rot caused by *F. solani* (isolate # 0110111067) on *Dracaena deremensis* (Table 29). Disease inoculation occurred after the first application, with the exception of Acibenzolar where inoculation occurred after the second application. Disease ratings were obtained weekly from 9/21 to 12/19, and plant marketability rating obtained with the help of a local grower on 12/19. Disease severity was moderate to severe throughout the trial. Disease levels for all fungicide treatments were statistically lower than the inoculated control. Though not statistically different from the other fungicide treatments, disease severity (stem rot) ratings for Heritage and 3336 were lowest and 100% of the canes treated with these products were in marketable condition. Also, disease severity from BW 240, CG100, Medallion, Palladium, and the higher rates of Tourney and Trinity were comparable to that from the non-inoculated check. No phytotoxicity was observed for any treatment.

**Table 28. Efficacy on Stem Root Rot Caused by *Fusarium solani* on *Dracaena deremensis* 'Janet Craig', Palmateer FL, 2011.**

Treatment	Rate Per 100 Gal	Application Method	Applic. Dates <sup>y</sup>	Applic. Interval	% Disease Severity <sup>x</sup>	Plant Marketability <sup>z</sup>
3336 F (thiophanate-methyl)	12 fl oz	Sprenc	1, 4	21 Days	5 d	5
Acibenzolar-s-methyl	0.125	Drench	1, 4	21 Days	37 b	2
	0.25 oz	Drench	1, 4	21 Days	37 b	2
BW240 WP ( <i>Trichoderma harzianum</i> & <i>T. virens</i> )	6 oz	Sprenc	2, 4	14 Days	12 cd	4
CG100 (caprylic acid)	0.6 pints	Sprenc	1, 3, 5, 6	14 Days	25 bc	3
Compass 50WDG (trifloxystrobin)	2 oz	Sprenc	1, 3, 5, 6	14 Days	36 b	2
Heritage 50WG (azoxystrobin)	4 oz	Sprenc	1, 5	28 Days	3 d	5
Medallion (fludioxonil)	2 oz	Sprenc	1, 4	21 Days	23 bc	3
Palladium 62.5WG (cyprodinil+fludioxonil)	4 oz	Sprenc	1, 3, 5, 6	14 Days	17 cd	4
Tourney 50 WDG (metconazole)	1 oz	Sprenc	1, 3, 5, 6	14 Days	18 cd	4
	2 oz	Sprenc	1, 3, 5, 6	14 Days	28 bc	3
Trinity 2 SC (triticonazole)	6 oz	Sprenc	1, 3, 5, 6	14 Days	18 cd	4
	8 oz	Sprenc	1, 3, 5, 6	14 Days	26 bc	3
Untreated non-inoculated	-	-	-	-	13 cd	4
Untreated inoculated	-	-	-	-	69 a	0

<sup>x</sup> Means followed by the same letter do not differ significantly based on based on Student Newman Keuls Test, (P=0.05).

<sup>y</sup> Dates: 1 = 9/1/11, 2 = 9/6/11, 3 = 9/15/11, 4 = 9/20/11, 5 = 9/29/11, 6 = 10/13/11.

<sup>z</sup> Rating of 0-5 where 0= not marketable, 5 = best marketability.

In 2002, Kirk conducted a field trial to determine efficacy of BAS 500, Bravo Zn, Gem and Quadris for control of root rot caused by *Fusarium solani* f. sp. *phaseoli* on snap bean (*Phaseolus vulgaris*). All plots were inoculated and in-furrow applications of fungicides were made over the seed at planting. No treatments were significantly different from each other as measured by root rot index at the first harvest (32 DAP) but at the second harvest (72 DAP) all treatments had a significantly lower root rot index value than the untreated control (Table 18). No treatments were significantly different from each other in plant emergence and marketable yield. No phytotoxicity was observed from any treatment.

**Table 29. \* Efficacy on Fusarium Root Rot (*Fusarium solani* f. sp. *phaseoli*) on Snap Bean (*Phaseolus vulgaris*) ‘Hi-Style’, Kirk, MI, 2002.**

Treatment <sup>y</sup>	Rate Per 1000 ft	% Emergence <sup>x</sup>	Root Rot Index <sup>y</sup>		Marketable Yield (Lb/A)
			35 DAT	72 DAT	
BAS 500 2.09EC (pyraclostrobin)	0.77 pt	98.8 a	13.8 a	48.8 bcde	1104 a
Bravo ZN 6SC (chlorothalonil)	1.5 pt	96.3 a	21.3 a	55.0 bc	981 a
Gem 4EC (trifloxystrobin)	0.8 pt	96.3 a	16.9 a	52.5 bcde	1023 a
Quadris 2SC (azoxystrobin)	0.8 pt	98.8 a	16.9 a	51.3 bcde	1056 a
Untreated inoculated	-	96.9 a	30.0 a	78.8 a	1008 a

\* Not an IR-4 Experiment: F&N Tests Vol 58: ST007. Not all products tested included in table.

<sup>x</sup> Means followed by same letter do not differ significantly based on Tukey Multiple Comparison (P = 0.05).

<sup>y</sup> Root rot index calculated by counting the number of roots from a sample of 10 plants falling onto class 0 = no visible root rot; 1 = 1 - 10% girdling of tap root; 2 = 11 - 20% girdling of tap root; 3 = 21 - 50% girdling of tap root; 4 = 51 - 100% girdling of tap root. The number in each class is multiplied by the class number and summed. The sum is multiplied by a constant to express as a percentage. Indices of 0 - 50 cover the range 0 - 20% girdling; 51 - 75 cover the range 21 - 50% girdling and > 75 cover the range 50 - 100% girdling..

## **Efficacy Summary by Product/Active Ingredient**

A brief efficacy summary for select products is given below, with a reminder that there are very limited data available to draw definitive conclusions for each product/pest species. Products were selected based on interest in these products for testing in 2010 and 2011 Fusarium efficacy projects.

**Acibenzolar.** In greenhouse trials, Acibenzolar applied as drench provided good efficacy against Fusarium wilt (*F. oxysporum*) on lisianthus, promising efficacy against Fusarium root rot (*F. oxysporum*) on larkspur, and mediocre efficacy on stem root rot (*F. solani*) on *Dracaena deremensis*. When applied as corm dip or foliar spray on corn flag, Acibenzolar and Acibenzolar + Heritage provided little efficacy against stem based necrosis (*F. oxysporum*) in two trials. On Douglas fir, drench or a combination of seed soak and drench applications provided no efficacy against damping-off and root rot (*Fusarium commune*). Actigard + Heritage applied as soil incorporation provided virtually no and mediocre efficacy in single trials on Douglas fir and bulbous iris. On vegetables, Actigard applied as drench provided poor to excellent efficacy against *Fusarium oxysporum* f. sp. *niveum* on watermelon in 5 greenhouse trials. In 6 field trials, Actigard provided good control when applied as drench to the soil immediately after transplanting, but variable control when applied through drip irrigation.

**Azoxystrobin.** Heritage applied as drench provided excellent efficacy against Fusarium crown and stem rot (*F. avenaceum*) on lisianthus, and on stem root rot (*F. solani*) on *Dracaena deremensis* in 2 greenhouse trials; however efficacy against severe Fusarium wilt (*F. oxysporum*) pressure was poor in two other trials. Single trials on Douglas fir showed drench or soil incorporation providing no efficacy against damping-off and root rot (*Fusarium commune*); mediocre efficacy was obtained with soil incorporation on bulbous iris. Used as a tuber soak, it did not significantly reduce disease severity of Fusarium tuber rot (*F. solani*) on caladium in one field trial. Quadris applied as drench provided good to excellent control of *Fusarium oxysporum* f. sp. *niveum* on watermelon in three greenhouse trials; however it provided no to poor control in 5 field trials.

**Boscalid + Pyraclostrobin.** In a greenhouse trial, Pageant applied as drench provided mediocre efficacy against Fusarium root rot (*F. oxysporum*) on larkspur. It provided promising and good efficacy against damping-off and root rot (*Fusarium commune*) applied as drench, and no efficacy as foliar, in 2 trials on Douglas fir.

**Caprylic Acid.** In greenhouse trials, CG100 applied as drench provided promising efficacy against stem root rot (*F. solani*) on *Dracaena deremensis*, poor efficacy against Fusarium root rot (*F. oxysporum*) on larkspur and against Fusarium wilt (*F. oxysporum*) on lisianthus. It provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag. Similarly, it provided no to poor efficacy against damping-off and root rot (*Fusarium commune*) was obtained on Douglas fir and bulbous iris when applied as drench or soil incorporation.

**Cyprodinil + Fludioxonil.** In greenhouse trials, Palladium applied as drench provided good efficacy against stem root rot (*F. solani*) on *Dracaena deremensis*, mediocre efficacy against Fusarium wilt (*F. oxysporum*) on lisianthus. It provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag. Similarly, it provided no efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir and bulbous iris when applied as drench or soil incorporation.

**Fludioxonil.** Medallion applied as drench provided excellent efficacy against severe Fusarium wilt (*F. oxysporum*) pressure on lisianthus in two greenhouse trials; however, it provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag in two trials. It also



provided poor efficacy against Fusarium crown and stem rot (*F. avenaceum*) in a lisianthus trial. Promising efficacy on stem root rot (*F. solani*) was obtained in *Dracaena deremensis*. On larkspur, it provided good efficacy against Fusarium root rot (*F. oxysporum*), but no efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir in 3 trials. Cannonball applied as drench provided no to poor efficacy against *Fusarium oxysporum* f. sp. *niveum* on watermelon in 5 greenhouse trials.

**Fludioxonil + Mefenoxam.** Hurricane applied as drench provided excellent efficacy against a severe Fusarium wilt (*F. oxysporum*) pressure on lisianthus in a greenhouse trial. In 3 trials, it provided good efficacy against Fusarium root rot (*F. oxysporum*) on larkspur, but no efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir. When applied as a tuber soak or as a spray at planting, it significantly reduced the severity of Fusarium tuber rot rot (*F. solani*) on caladium in a field trial.

**Fluoxastrobin.** Disarm applied as corm dip or foliar spray provided significant, but poor, efficacy against stem based necrosis (*F. oxysporum*) on corn flag in two trials.

**Metconazole.** In greenhouse trials, Tourney applied as drench provided good to excellent efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir; when applied as soil incorporation, it provided good efficacy on bulbous iris but no efficacy on Douglas fir. It provided good efficacy against Fusarium wilt (*F. oxysporum*) on lisianthus, and against stem root rot (*F. solani*) on *Dracaena deremensis*, but poor efficacy against Fusarium root rot (*F. oxysporum*) on larkspur. Tourney provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag in two trials. V-10116 or Metconazole applied as drench provided good to excellent control of *Fusarium oxysporum* f. sp. *niveum* on watermelon in 3 greenhouse trials; however poor efficacy was observed in 2 field trials.

**Prothioconazole.** Proline applied as drench provided good to excellent efficacy against *Fusarium oxysporum* f. sp. *niveum* on watermelon in 3 greenhouse trials. It also provided good control in 6 field trials either applied as drench to the soil immediately after transplanting or through drip irrigation.

**Pyraclostrobin.** In a greenhouse trial, Insignia provided no to poor efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir. Cabrio applied as drench provided good efficacy against *Fusarium oxysporum* f. sp. *niveum* on watermelon in one trial but no efficacy in 2 other greenhouse trials. BAS 500 applied in-furrow significantly reduced Fusarium root rot caused by *Fusarium solani* f. sp. *phaseoli* on snap bean in one field trial.

**SP2169.** In greenhouse trials, SP2169 applied as drench provided good efficacy against Fusarium wilt (*Fusarium oxysporum*) on lisianthus but no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag in two trials. Similarly, it provided no efficacy against damping-off and root rot (*F. commune*) on Douglas fir and bulbous iris when applied as drench or soil incorporation.

**Thiamethoxam.** In two greenhouse trials, Flagship 0.22G applied as soil incorporation provided good efficacy against damping-off and root rot (*F. commune*) on Douglas fir and bulbous iris; however Flagship 25WG provided no efficacy. In the same trials, Flagship + Heritage generally provided slight improvement in efficacy compared to Flagship or Heritage alone.

**Tebuconazole.** In a greenhouse trial, Torque applied as drench provided excellent efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir. However, Torque applied as soil incorporation in two other trials provided poor efficacy against damping-off and root rot on bulbous iris and no efficacy on Douglas fir.

**Thiophanate methyl.** In greenhouse trials, 3336 applied as drench provided excellent efficacy against stem root rot (*F. solani*) on *Dracaena deremensis*, and good efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir and against Fusarium root rot (*F. oxysporum*) on larkspur. It provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag in two trials. It provided poor efficacy against Fusarium crown and stem rot (*F. avenaceum*) and poor to mediocre efficacy against Fusarium wilt (*F. oxysporum*) on lisianthus in 3 greenhouse trials. Used as a tuber soak, it did not significantly reduce disease severity of Fusarium tuber rot (*F. solani*) on caladium in one field trial. On vegetables, Topsin applied as drench provided excellent efficacy against *Fusarium oxysporum* f. sp. *niveum* on watermelon in 4 of 5 greenhouse trials. In 4 field trials, Topsin provided good control when applied as drench to the soil immediately after transplanting, but variable control when applied through drip irrigation.

**Trichoderma harzianum & T. virens.** In greenhouse trials, BW240 applied as drench or soil incorporation provided good efficacy against stem root rot (*F. solani*) on *Dracaena deremensis*, poor efficacy against Fusarium root rot (*F. oxysporum*) on larkspur, and no efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir and bulbous iris, and against Fusarium wilt (*F. oxysporum*) on lisianthus. Also it provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as corm dip or foliar spray on corn flag in two trials.

**Trifloxystrobin.** In greenhouse trials, Compass applied as drench provided mediocre efficacy on stem root rot (*F. solani*) on *Dracaena deremensis*, no and promising activity against damping-off and root rot (*Fusarium commune*) on Douglas fir, and poor efficacy against a severe Fusarium wilt (*F. oxysporum*) pressure on lisianthus. Used as a tuber soak, it did not significantly reduce disease severity of Fusarium tuber rot (*F. solani*) on caladium in one field trial. Gem applied in-furrow significantly reduced Fusarium root rot caused by *Fusarium solani* f. sp. *phaseoli* on snap bean in one field trial.

**Triflumizole.** Terraguard applied as drench provided excellent efficacy against a severe Fusarium wilt (*F. oxysporum*) pressure on lisianthus in a greenhouse trial.

**Triticonazole.** In greenhouse trials, Trinity applied as drench provided good efficacy against stem root rot (*F. solani*) on *Dracaena deremensis*, but no and promising efficacy against damping-off and root rot (*Fusarium commune*) on Douglas fir. Efficacy was promising against Fusarium wilt (*F. oxysporum*) on lisianthus, but poor against Fusarium root rot (*F. oxysporum*) on larkspur in other trials. Trinity applied as soil incorporation provided good efficacy against damping-off and root rot (*F. commune*) on bulbous iris but no efficacy on Douglas fir. Also it provided no efficacy against stem based necrosis (*F. oxysporum*) when applied as foliar spray on corn flag. This active ingredient applied as drench provided good control of *Fusarium oxysporum* f. sp. *niveum* on watermelon in one greenhouse trial but no control in 2 other trials.

### **Phytotoxicity**

No phytotoxicity was observed with the products listed above with the exception of acibenzolar on watermelon (stunting, leaf yellowing) and metconazole on watermelon (stunting, dark green leaves) in some trials. Significant stunting was also observed with metconazole on lisianthus and SP2169 on Douglas fir.

**Table 30. Summary of product efficacy by pathogen and crop.**

Note: Table entries are sorted by product, pathogen Latin name, and then by crop Latin name. Only those IR-4 trials received by 5/5/2013 are included in the table below.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31280	3336 F (Thiophanate-methyl)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Significantly increased healthy seedlings with 16 fl oz per 100 gal; comparable to non-inoculated Check.
30475	3336 F (Thiophanate-methyl)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Excellent control at 4 lb per 100 gal applied once; almost comparable to non-inoculated check; very minor, transient leaf phytotoxicity.
31380	3336 F (Thiophanate-methyl)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Significantly reduced a moderate to severe disease severity with 12 oz per 100 gal applied twice; comparable to non-inoculated check.
31378	3336 F (Thiophanate-methyl)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 16 fl oz per 100 gal.
31378	3336 F (Thiophanate-methyl)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 16 fl oz per 100 gal.
29745	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	No control of a high disease pressure with 0.125 and 0.25 oz per 100 gal.
29745	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Seed soak and drench treatments.	High disease pressure. Did not increase healthy seedlings with 0.25 oz per 100 gal.
30468	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Significantly reduced leaf and root necrosis at 0.25 oz per 100 gal applied once; inferior to Medallion; very minor, transient leaf phytotoxicity.
30244	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Drench	Significantly reduced a moderate to severe disease severity with 0.125 and 0.25 oz per 100 gal applied twice; inferior to non-inoculated check.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31369	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Poor reduction of stem base necrosis incidence with 0.125 and 0.25 oz per 100 gal.
31369	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 0.5 and 0.75 oz per 100 gal.
30191	Acibenzolar-S-methyl (Acibenzolar-S-methyl)	Fusarium sp. (Fusarium sp.)	Lisianthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Significantly reduced F. oxysporum severity with 0.125 and 0.25 oz per 100 gal applied every 21 days; comparable to non-inoculated check.
29747	CG100 (Caprylic acid)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Little reduction of high disease pressure with 0.6 pt pe 100 gal.
29747	CG100 (Caprylic acid)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Significantly increased healthy seedlings with 0.6 pt per 100 gal; inferior to non-inoculated Check.
29747	CG100 (Caprylic acid)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 26 ml per 1000 sq ft.
30470	CG100 (Caprylic acid)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Significantly reduced leaf and root necrosis at 0.8 pt per 100 gal applied once; inferior to Medallion; very minor, transient leaf phytotoxicity.
30246	CG100 (Caprylic acid)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Some reduction with a moderate to severe disease severity with 0.8 pt per 100 gal applied 4 times; slightly inferior to non-inoculated check.
31371	CG100 (Caprylic acid)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 38.4 fl oz per 100 gal.
31371	CG100 (Caprylic acid)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 38.4 fl oz per 100 gal.
31391	CG100 (Caprylic acid)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	No significant control with 26 ml per 1000 sq ft.
30193	CG100 (Caprylic acid)	Fusarium sp. (Fusarium sp.)	Lisianthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Did not significantly reduce F. oxysporum severity with 0.6 pt per 100 gal applied every 14 days.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30000	Compass 0 50WDG (Trifloxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Some control of a high disease pressure with 0.5 oz per 100 gal.
30000	Compass 0 50WDG (Trifloxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Some control with 0.5 oz per 100 gal.
31382	Compass 0 50WDG (Trifloxystrobin)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Some reduction with moderate to severe disease severity with 2 oz per 100 gal applied 4 times; inferior to non-inoculated check.
31377	Disarm 480SC (Fluoxastrobin)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Significant, but mediocre, reduction of stem base necrosis incidence with 4 fl oz per 100 gal.
31377	Disarm 480SC (Fluoxastrobin)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 4 fl oz per 100 gal.
31384	Flagship 0.22G (Thiamethoxam)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 44 oz per 1000 sq ft.
31401	Flagship 0.22G (Thiamethoxam)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	Significantly increased total marketable and 18" tall flowers with 44 oz per 1000 sq ft.
31385	Flagship 25WG (Thiamethoxam)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 0.55 oz per 1000 sq ft.
29753	Heritage (Azoxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Poor to some control of a high disease pressure with 1.8 and 4.0 oz per 100 gal.
29753	Heritage (Azoxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Did not significantly increase healthy seedlings with 1.8 oz per 100 gal.
29753	Heritage (Azoxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 0.4 oz per 1000 sq ft.
31379	Heritage (Azoxystrobin)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Significantly reduced a moderate to severe disease severity with 4 oz per 100 gal applied twice; comparable to non-inoculated check.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31400	Heritage (Azoxystrobin)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	No significant control with 0.4 oz per 1000 sq ft.
30001	Hurricane (fludioxonil + mefonaxam)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	No control of a high disease pressure with 0.75 oz per 100 gal.
30001	Hurricane (fludioxonil + mefonaxam)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Did not significantly increase healthy seedlings with 0.75 oz per 100 gal.
30473	Hurricane (fludioxonil + mefonaxam)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Significantly reduced leaf and root necrosis at 12 oz per 100 gal applied once; comparable to Medallion; very minor, transient leaf phytotoxicity.
31281	Insignia SC (Pyraclostrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Significantly increased healthy seedlings with 6.1, but not with 3.1, fl oz per 100 gal; inferior to non-inoculated Check.
30474	Medallion (Fludioxonil)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Excellent control at 2 fl oz per 100 gal applied once; almost comparable to non-inoculated check; very minor, transient leaf phytotoxicity.
31381	Medallion (Fludioxonil)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Drench	Some reduction with a moderate to severe disease severity with 2 oz per 100 gal applied twice; inferior to non-inoculated check.
31376	Medallion (Fludioxonil)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 8 oz per 100 gal.
31376	Medallion (Fludioxonil)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 4 oz per 100 gal.
29749	Pageant 38WG (Boscalid + Pyraclostrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Good control of a high disease pressure with 12 oz per 100 gal.
29749	Pageant 38WG (Boscalid + Pyraclostrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Significantly increased healthy seedlings with 12 oz per 100 gal; comparable to non-inoculated Check.
29749	Pageant 38WG (Boscalid + Pyraclostrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Foliar	Poor control of a high disease pressure with 12 oz per 100 gal.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30476	Pageant 38WG (Boscalid + Pyraclostrobin)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Significantly reduced leaf and root necrosis at 12 oz per 100 gal applied twice; inferior to Medallion; very minor, transient leaf phytotoxicity.
29750	Palladium (Cyprodinil + fludioxanil)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	No control of a high disease pressure with 6 oz per 100 gal.
29750	Palladium (Cyprodinil + fludioxanil)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Did not significantly increase healthy seedlings with 4 oz per 100 gal.
29750	Palladium (Cyprodinil + fludioxanil)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not increase number of healthy seedlings with 1 oz per 1000 sq ft.
30247	Palladium (Cyprodinil + fludioxanil)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Spreng	Significantly reduced a moderate to severe disease severity with 6 oz per 100 gal applied 4 times; comparable to non-inoculated check.
31372	Palladium (Cyprodinil + fludioxanil)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 6 fl oz per 100 gal.
31372	Palladium (Cyprodinil + fludioxanil)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 6 fl oz per 100 gal.
31392	Palladium (Cyprodinil + fludioxanil)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	No significant control with 1 oz per 1000 sq ft.
30194	Palladium (Cyprodinil + fludioxanil)	Fusarium sp. (Fusarium sp.)	Lisianthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Did not significantly reduce F. oxysporum severity with 6 oz per 100 gal applied every 14 days.
29744	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	No control of a high disease pressure with 8 oz per 100 gal.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
29744	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Potting mix and seed treatments	High disease pressure. No control with 6 oz per 100 gal potting mix and seed treatments.
29744	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 10 per 1000 sq ft.
30469	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Root dip, Drench	Significantly reduced leaf and root necrosis at 6, then 3 oz per 100 gal; inferior to Medallion; very minor, transient leaf phytotoxicity.
30245	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Spreng	Significantly reduced a moderate to severe disease severity with 6 oz per 100 gal applied twice; comparable to non-inoculated check.
31370	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 80 oz per 100 gal.
31370	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 8 oz per 100 gal.
31390	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	No significant control with 10 oz per 1000 sq ft.



PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30192	RootShield Plus WP (aka BW240) (Trichoderma harzianum T-22 + Trichoderma virens G-41)	Fusarium sp. (Fusarium sp.)	Lisianthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Did not significantly reduce F. oxysporum severity with 6 oz, then 3 oz per 100 gal.
30222	SP2169 (SP2169)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Did not significantly increase healthy seedlings with 12.3 oz per 100 gal; significant phytotoxicity.
30222	SP2169 (SP2169)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 8.35 ml per 1000 sq ft.
31373	SP2169 (SP2169)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 12.3 fl oz per 100 gal.
31373	SP2169 (SP2169)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 12.3 fl oz per 100 gal.
31393	SP2169 (SP2169)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	No significant control with 8.35 ml per 1000 sq ft.
30195	SP2169 (SP2169)	Fusarium sp. (Fusarium sp.)	Lisianthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Significantly reduced F. oxysporum severity with 12.3 oz per 100 gal applied every 14 days; comparable to non-inoculated check.
31383	Tank Mix: Acibenzolar + Heritage (Acibenzolar + Azoxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not increase number of healthy seedlings with 0.08 g + 1 oz per 1000 sq ft.
31546	Tank Mix: Acibenzolar + Heritage (Acibenzolar + Azoxystrobin)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Reduction of stem base necrosis incidence mediocre with 0.125 + 0.45, poor with 0.25 + 0.9 and 0.5 + 1.8, oz per 100 gal.
31546	Tank Mix: Acibenzolar + Heritage (Acibenzolar + Azoxystrobin)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not reduce stem base necrosis incidence with 0.5 + 1.8 and 0.75 + 2.7 oz per 100 gal.
31397	Tank Mix: Acibenzolar + Heritage (Acibenzolar + Azoxystrobin)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	Significantly increased 18" tall flowers with 0.08 g + 1 oz per 1000 sq ft.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
31387	Tank Mix: Flagship 25WG + Heritage (Thiamethoxam + azoxystrobin)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Only treatment that significantly increased number of healthy seedlings with 0.4 oz + 0.4 oz per 1000 sq ft.
31399	Tank Mix: Flagship 25WG + Heritage (Thiamethoxam + azoxystrobin)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	Significantly increased 18" tall flowers with 0.4 oz + 0.4 oz per 1000 sq ft.
31282	Torque 3.6SC (Tebuconazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not increase number of healthy seedlings with 20.4 ml per 1000 sq ft.
29751	Tourney 50WDG (Metconazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Excellent control of a high disease pressure with 1 and 2 oz per 100 gal; best treatment.
29751	Tourney 50WDG (Metconazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Significantly increased healthy seedlings with 1 and 2 oz per 100 gal; comparable to non-inoculated Check.
29751	Tourney 50WDG (Metconazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 2.8 and 5.6 g per 1000 sq ft.
30471	Tourney 50WDG (Metconazole)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Did not significantly reduce leaf and root necrosis at 2 fl oz per 100 gal applied once; very minor, transient leaf phytotoxicity.
30249	Tourney 50WDG (Metconazole)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Significantly reduced a moderate to severe disease severity with 1 and 2 oz per 100 gal applied 4 times; slightly inferior to non-inoculated check.
31374	Tourney 50WDG (Metconazole)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Bulb Dip	Did not significantly reduce stem base necrosis incidence with 2 and 4 fl oz per 100 gal.
31374	Tourney 50WDG (Metconazole)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 1 and 2 fl oz per 100 gal.
31394	Tourney 50WDG (Metconazole)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	Significantly increased total marketable and 18" tall flowers, and decrease culls with 5.6 g per 1000 sq ft; lower rate less effective.

PR#	Product (Active ingredient)	Target	Crop	Production Site	Researcher	State	Year	Application Type	Results
30196	Tourney 50WDG (Metconazole)	Fusarium sp. (Fusarium sp.)	Lisanthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Did not significantly reduce F. oxysporum severity with 1 and 2 oz per 100 gal applied every 14 days; severe stunting.
29752	Trinity 2SC (Triticonazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2010	Drench	Great control (comparable to non inoculated) of a high disease pressure with 6 and 8 fl oz per 100 gal.
29752	Trinity 2SC (Triticonazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2011	Drench	High disease pressure. Did not significantly increase healthy seedlings with 6 and 8 fl oz per 100 gal; slight phyto at the higher rate.
29752	Trinity 2SC (Triticonazole)	Fusarium commune (Fusarium commune)	Fir, Douglas (Pseudotsuga menziesii)	Seedbed	Chastagner	WA	2012	Soil incorporation	Did not significantly increase number of healthy seedlings with 5.43 and 8.14 ml per 1000 sq ft.
30472	Trinity 2SC (Triticonazole)	Fusarium oxysporum (Fusarium oxysporum)	Larkspur (Delphinium sp.)	Greenhouse	Kirk	MI	2010	Drench	Significantly reduced leaf and root necrosis at 11.8 oz per 100 gal applied as curative twice; inferior to Medallion; very minor, transient leaf phytotoxicity.
30250	Trinity 2SC (Triticonazole)	Fusarium solani (Fusarium solani)	Dracaena deremensis (Dracaena deremensis) 'Janet Craig'	Greenhouse	Palmeteer	FL	2011	Sprenc	Significantly reduced a moderate to severe disease severity with 6 and 8 oz per 100 gal applied 4 times; slightly inferior to non-inoculated check.
31375	Trinity 2SC (Triticonazole)	Fusarium sp. (Fusarium sp.)	Corn Flag, Sword Lily (Gladiolus sp.) 'Sun-Kissed'	Field In-Ground	Kirk	MI	2012	Foliar	Did not significantly reduce stem base necrosis incidence with 8 and 12 fl oz per 100 gal.
31395	Trinity 2SC (Triticonazole)	Fusarium sp. (Fusarium sp.)	Iris, Dutch (Iris x hollandica) 'Blue Diamond'	Field Container	Chastagner	WA	2012	Soil incorporation	Significantly increased total marketable and 18" tall flowers, and decrease culls with 8.14 ml per 1000 sq ft; best treatment; lower rate less effective.
30197	Trinity 2SC (Triticonazole)	Fusarium sp. (Fusarium sp.)	Lisanthus (Lisianthus sp.) Eustoma grandiflora	Greenhouse	Chase	CA	2011	Drench	Did not significantly reduce F. oxysporum severity with 6 and 8 oz per 100 gal applied every 14 days; significant stunting at the higher rate.

## Appendix 1: Contributing Researchers

Dr. Ann Chase	Chase Horticultural Research 8031 Mount Aukum Road Ste F P. O. Box 529 Mount Aukum, CA 95656-0529
Dr. Gary Chastagner	Washington State University Research and Extension Center 7612 Pioneer Way East Puyallup, WA 98371
Dr. Dan Egel	Purdue University Dept. of Botany and Plant Pathology Southwest Purdue Ag Center 4369 N. Purdue Rd. Vincennes, IN 47591
Dr. Kathryne. L. Everts	University of Maryland Lower Eastern Shore Research & Education Center 27664 Nanticoke Road Salisbury, MD 21801
Dr. William W. Kirk	Michigan State University Dept. of Plant Pathology 35 Plant Pathology Building East Lansing, MI 48824
Dr. D. B. Langston, Jr.	University of Georgia Dept. of Plant Pathology 4604 Research Way Tifton, GA 31793
Dr. Robert McGovern ( <i>retired</i> )	University of Florida Gulf Coast Research and Education Center 14625 CR 672 Wimauma, FL 33598
Dr. Aaron Palmateer	University of Florida Tropical Research & Education Center 18905 SW 280 Street Homestead, FL 33031

## **Appendix 2: Submitted Data Reports**

The IR-4 reports in this Appendix cover multiple PR numbers and are arranged alphabetically by the researchers' last names. Only those reports received by 5/5/2013 are included.

These reports can also be found at [www.rutgers.ir4.edu](http://www.rutgers.ir4.edu) by searching under the Fusarium Efficacy project.